

Yuh-Chin T Huang MD MHS

Department of Medicine
Duke University Medical Center
Durham, North Carolina

The authors have disclosed no conflicts of interest.

REFERENCES

1. Su CL, Chiang LL, Yang SH, Lin HI, Cheng KC, Huang YC, Wu CP. Preventive use of

noninvasive ventilation after extubation: a prospective, multicenter randomized controlled trial. *Respir Care* 2012;57(2):204-210.

2. Ferrer M, Esquinas A, Arancibia F, Bauer TT, Gonzalez G, Carrillo A, et al. Noninvasive ventilation during persistent weaning failure: a randomized controlled trial. *Am J Respir Crit Care Med* 2003;168(1):70-76.
3. Ferrer M, Valencia M, Nicolas JM, Bernadich O, Badia JR, Torres A. Early non-

invasive ventilation averts extubation failure in patients at risk: a randomized trial. *Am J Respir Crit Care Med* 2006;173(2):164-170.

4. Nava S, Gregoretti C, Fanfulla F, Squadrone E, Grassi M, Carlucci A, et al. Noninvasive ventilation to prevent respiratory failure after extubation in high-risk patients. *Crit Care Med* 2005;33(11):2465-2470.

DOI: 10.4187/respcare.01969

Notice of Retraction

At the request of the author, "Comparing performance of 3 oscillating positive expiratory pressure devices at similar amplitude and frequencies of oscillations on displacement of mucus inside trachea during cough" has been withdrawn. The paper has not been published in print, but was published electronically ahead of print on March 13, 2012. DOI: 10.4187/respcare.01631.

DOI: 10.4187/respcare.02033