

Quantitative Computed Tomography and Response to Pronation in COVID-19 ARDS

Francesco Zadek, Luca Berta, Giulia Zorzi, Stefania Ubiali, Amos Bonaiti, Giulia Tundo, Beatrice Brunoni, Francesco Marrazzo, Riccardo Giudici, Anna Rossi, Francesco Rizzetto, Davide Paolo Bernasconi, Angelo Vanzulli, Paola Enrica Colombo, Roberto Fumagalli, Alberto Torresin, and Thomas Langer

BACKGROUND: The use of prone position (PP) has been widespread during the COVID-19 pandemic. Whereas it has demonstrated benefits, including improved oxygenation and lung aeration, the factors influencing the response in terms of gas exchange to PP remain unclear. In particular, the association between baseline quantitative computed tomography (CT) scan results and gas exchange response to PP in invasively ventilated subjects with COVID-19 ARDS is unknown. The present study aimed to compare baseline quantitative CT results between subjects responding to PP in terms of oxygenation or CO₂ clearance and those who did not. **METHODS:** This was a single-center, retrospective observational study including critically ill, invasively ventilated subjects with COVID-19-related ARDS admitted to the ICUs of Niguarda Hospital between March 2020–November 2021. Blood gas samples were collected before and after PP. Subjects in whom the P_{aO₂}/F_{IO₂} increase was ≥ 20 mm Hg after PP were defined as oxygen responders. CO₂ responders were defined when the ventilatory ratio (VR) decreased during PP. Automated quantitative CT analyses were performed to obtain tissue mass and density of the lungs. **RESULTS:** One hundred twenty-five subjects were enrolled, of which 116 (93%) were O₂ responders and 51 (41%) CO₂ responders. No difference in quantitative CT characteristics and oxygen were observed between responders and non-responders (tissue mass 1,532 ± 396 g vs 1,654 ± 304 g, *P* = .28; density −544 ± 109 HU vs −562 ± 58 HU *P* = .42). Similar findings were observed when dividing the population according to CO₂ response (tissue mass 1,551 ± 412 g vs 1,534 ± 377 g, *P* = .89; density −545 ± 123 HU vs −546 ± 94 HU, *P* = .99). **CONCLUSIONS:** Most subjects with COVID-19-related ARDS improved their oxygenation at the first pronation cycle. The study suggests that baseline quantitative CT scan data were not associated with the response to PP in oxygenation or CO₂ in mechanically ventilated subjects with COVID-19-related ARDS. *Key words:* computed tomography; quantitative CT scan analysis; prone position; COVID-19 ARDS; coronavirus SARS; pulmonary gas exchange; lung compliance; ventilation/perfusion scan. [Respir Care 2024;0(0):1–●. © 2024 Daedalus Enterprises]

Introduction

Prone position (PP) has been extensively used during the COVID-19 pandemic in invasively ventilated subjects.¹ The benefits reported from the use of this position in classic ARDS were also confirmed in COVID-19-associated ARDS.² This strategy, requiring highly trained personnel and not devoid of possible complications,^{3–5} has thus been included in the guidelines for the treatment of moderate and severe COVID-19-associated ARDS.⁶ Indeed, whereas results from randomized controlled

trials in this specific population are lacking, placing subjects with COVID-19-associated ARDS in PP decreases alveolar collapse, hyperinflation, and improves the homogeneity of lung aeration and ventilation.^{2,7} Moreover, whereas not the primary target of PP, several studies reported a variable (ie, between 30–80%) improvement in oxygenation during PP of mechanically ventilated subjects with COVID-19-associated ARDS.^{1,2} However, it is currently unknown which factors contribute to and how to predict the response in terms of oxygenation in patients with ARDS placed in PP.

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

Chest computed tomography (CT) was broadly used in patients with COVID-19 to facilitate diagnosis and quantify the degree of disease extension.^{8,9} Several radiological patterns could be observed at different times throughout the disease course, showing diffuse lung alterations ranging from ground-glass opacities to parenchymal consolidations.^{10,11} In addition, quantitative CT results, cornerstones for the understanding of classic ARDS,¹² have been analyzed to investigate the pathophysiology of COVID-19-associated ARDS and the lung response to PP in selected groups of subjects.^{2,13} Previous studies have suggested that in supine position the amount of non-aerated lung tissue in the dependent lung regions was associated with more recruitable lung volume when PP was used,¹⁴⁻¹⁶ and recently, a relationship between the dorsal non-aerated tissue quantified at the CT scan and the gas exchange response to PP was recorded in classic ARDS.¹⁷ These studies, however, did not focus on the association between quantitative CT results and the oxygenation response to PP in subjects with COVID-19-related ARDS.

Recently, Raimondi et al¹⁸ studied awake, noninvasively ventilated subjects with COVID-19 and were not able to find any association between the distribution of CT lung lesions and the response in oxygenation to PP. Information regarding the association of baseline quantitative CT results and the response to PP in invasively ventilated patients with COVID-19-associated ARDS is currently lacking. We hypothesized that the quantitative CT results of scans performed prior to the first PP would differ significantly between responders and non-responders in terms of oxygenation and CO₂ clearance. The present retrospective study was conducted to test this hypothesis.

Methods

Study Design

This was a single-center, retrospective, observational study performed at the Grande Ospedale Metropolitano Niguarda in Milan, Italy. The retrospective access to clinical data was approved by the ethical committee Milano Area B (approval number: 593-06102020), and the need for informed consent from individual subjects was waived.

Drs Zadek, Bonaiti, Tundo, and Brunoni are affiliated with Department of Medicine and Surgery, University of Milan-Bicocca, Monza, Italy. Drs Berta, Colombo, and Torresin are affiliated with Department of Medical Physics, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy. Ms Zorzi is affiliated with Department of Medical Physics, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; and Department of Physics, INFN Milan Unit, Milan, Italy. Dr Ubiali is affiliated with Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy. Drs Marrazzo, Giudici, and Rossi are affiliated with Department of Anesthesia and Intensive Care Medicine, Niguarda Ca' Granda, Milan, Italy. Dr Rizzetto is affiliated with Department of Radiology, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy. Dr Bernasconi is affiliated with School of Medicine and Surgery, Bicocca Bioinformatics Biostatistics and Bioimaging Center - B4, University of Milano-Bicocca, Monza, Italy.

QUICK LOOK

Current knowledge

Chest computed tomography (CT) has been broadly used in COVID-19 pandemic to facilitate diagnosis and quantify the degree of disease extension. Recently, a relationship between the dorsal non-aerated tissue quantified at the CT scan and the gas exchange response to prone position (PP) was demonstrated in classic ARDS, but clinical studies have not confirmed these findings in COVID-19-related ARDS.

What this paper contributes to our knowledge

No relationship between dorsal non-aerated tissue quantified at the CT scan analysis and either oxygen or CO₂ response to pronation was found. Quantitative CT imaging should not be accounted for when deciding whether to use PP in intubated patients with COVID-19-related ARDS.

All patients admitted between March 1, 2020–November 30, 2021, to the COVID-19 ICUs were screened for eligibility. Inclusion criteria were as follows: (1) age > 18 y, (2) laboratory-confirmed SARS-CoV-2 infection, (3) ARDS diagnosis according to Berlin criteria at ICU admission,¹⁹ (4) tracheal intubation and invasive mechanical ventilation, (5) use of PP, and (6) performance of a chest CT scan within the 72 h prior the first PP. Exclusion criteria were missing clinical data regarding blood gas analysis performed during the first PP cycle. Intubated subjects with COVID-19-related ARDS were maintained sedated and paralyzed. Subjects were ventilated using a lung-protective ventilatory strategy: low tidal volume (V_T ; 6–8 mL/predicted body weight), medium-high levels (8–12 cm H₂O) of PEEP, breathing frequency between 15–25 breaths/min, maintaining a plateau pressure < 28 cm H₂O and a driving pressure < 12 cm H₂O, and with a target S_{pO_2} of 92–95%. A P_{aO_2}/F_{IO_2} < 100 mm Hg was used as a criterion for PP.²⁰

Dr Vanzulli is affiliated with Department of Radiology, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; and Department of Oncology and Hemato-Oncology, Università degli Studi di Milano, Milan, Italy. Drs Fumagalli and Langer are affiliated with Department of Medicine and Surgery, University of Milan-Bicocca, Monza, Italy; and Department of Anesthesia and Intensive Care Medicine, Niguarda Ca' Granda, Milan, Italy.

The authors have disclosed no conflicts of interest.

Correspondence: Thomas Langer MD, Department of Medicine and Surgery, University of Milan-Bicocca, Monza, Italy. E-mail: Thomas.Langer@unimib.it.

DOI: 10.4187/respcare.11625

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

Clinical management, including the decision to use PP and perform a chest CT scan, was at the discretion of the attending physicians. The final date of follow-up for subject outcomes was July 14, 2022. For study purposes linked to the regional research network,²¹ an extensive set of information was prospectively recorded from the day of ICU admission on an electronic report form (REDCap electronic data capture tools). This information included anthropometric and clinical data, severity scores, vital signs, laboratory tests, radiological information, ICU and hospital length of stay (LOS), and ICU and hospital survival.

To assess the physiologic effects of pronation, subjects' ventilatory settings were prospectively recorded at 3 different time points: (1) within 2 h before the pronation (baseline); (2) during the last 4 h of the pronation cycle (prone); and (3) within 2–5 h after turning the subjects back to the supine position (supine). At each time point, end-inspiratory and end-expiratory airway occlusion maneuvers were performed to calculate driving pressure and respiratory system compliance (C_{RS}).²² At the same time points, arterial blood gases were drawn to calculate the P_{aO_2}/F_{IO_2} and the ventilatory ratio (VR).^{22,23}

Definitions

Subjects were defined as oxygen responders to PP according to 2 different definitions previously applied in the literature: (1) the P_{aO_2}/F_{IO_2} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position;¹ similarly, oxygen non-responders were defined as those subjects in whom this condition was not satisfied; and (2) the median P_{aO_2}/F_{IO_2} increase observed during prone ventilation was used as cutoff, defining responders with a P_{aO_2}/F_{IO_2} higher than the median value and non-responders with a P_{aO_2}/F_{IO_2} below.²⁴ The change in the VR was used to define the response in terms of CO_2 clearance. Subjects were defined as CO_2 responders when the VR decreased during pronation as compared to supine, while CO_2 non-responder when the VR increased or did not change.

CT Scan Acquisition and Image Analysis

All CT images were acquired on 4 scanners of a single vendor (Siemens, Munich, Germany) and with the same acquisition protocol for chest examinations, employing an automatic exposure control and an automatic selection of the tube voltage and sharp reconstruction algorithm.

All CT series were exported from the picture archiving communication system to a dedicated workstation for automatic image analysis. A dedicated processing software developed in Python language was used, as previously described.^{25–27} Briefly, (1) the pipeline rescales CT images to a slice thickness of 3 mm; (2) performs, for each slice, automatic segmentation of the left and right lungs; and (3)

calculates the relative distribution of Hounsfield units (HU) of the segmented regions of interest.

In this work, the following metrics were considered: volume (V_{lung} [mL]), Hounsfield unit related to the lung density (ρ [HU]), and mass (m [g]). The volume and the density were calculated, respectively, as the number of voxels multiplied by the physical voxel dimension and their average HU value of the selected region of interest. The mass was calculated using the following formula:

Lung tissue mass =

$$\rho \cdot V_{lung} \times (1,000 - \text{Hounsfield unit})/1,000$$

For each CT image, all these metrics were calculated for both lungs, obtained as the sum of the segmentations of the right and left lungs, and in 4 different density regions according to classical aeration thresholds:¹² hyperinflated lung [–1,000 to –900] HU, well-aerated lung [–900 to –500] HU, poorly aerated lung [–500 to –100] HU, and non-aerated lung [–100 to +100] HU.²⁸

Furthermore, a geometric subdivision of the entire (both lungs) region of interest was performed. The masks were divided into 10 different regions equally spaced along the sternovertebral axis, and for each subregion, the previous metrics were calculated.

Statistical Analysis

No sample size calculation was performed a priori, and the sample size is equal to the number of patients treated in our hospital during the study period. Comparison between continuous variables was performed via Student *t* test using Welch correction for unequal variance, Mann-Whitney rank-sum test, analysis of variance, or Kruskal-Wallis test as appropriate. Differences between categorical variables were assessed using the chi-square or Fisher exact test. The continuous relationship between quantitative variables was investigated using linear regression. Data were expressed as mean \pm SD or median and interquartile range. Statistical significance was defined as $P < .050$. Analyses were performed with Stata statistical software (release 16, StataCorp, College Station, Texas), and graphs were drawn using SigmaPlot v.12.0 (Systat Software, San Jose, California). The Standards of Reporting of Observational Studies in Epidemiology checklist for observational studies was used.

Results

During the study period, 466 patients with COVID-19 were admitted to the ICU (Fig. 1). One hundred twenty-five subjects with a median Simplified Acute Physiology Score II score at ICU admission of 38 [33–43] were enrolled in

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS




Fig. 1. Study flow chart. PP = prone position; CT = computed tomography; ABG = arterial blood gas.

the study. Baseline demographic characteristics are summarized in Table 1.

Oxygen Response to Pronation and Quantitative CT Scan Parameters

According to the P_{aO_2}/F_{IO_2} increased by ≥ 20 mm Hg definition, 116 subjects (93%) were O_2 responders, while

9 (7%) were non-responders. O_2 responders had a higher body mass index ($P = .009$) and prevalence of hypertension ($P = .001$) compared to nonresponders. The use of noninvasive respiratory support prior to intubation (72% vs 100%, $P = .063$), its duration (1 [0–3] d vs 1 [1–4] d, $P = .08$), and the use of awake PP prior to intubation (37% vs 44% $P = .44$) were similar between O_2 responders and non-responders. No difference in ARDS severity, ventilatory settings, and blood gas parameters was recorded (Table 2).

O_2 responders were characterized by higher baseline C_{RS} (42 ± 115 mL/cm H_2O vs 32 ± 5 mL/cm H_2O , $P \leq .001$). The length of the first pronation performed in the ICU was similar in O_2 responders and non-responders (21 [18–24] h vs 24 [22–32] h, $P = .08$).

During the first PP, O_2 responders improved, as per the definition, the P_{aO_2}/F_{IO_2} . Moreover, arterial pH and C_{RS} increased, while the VR did not change significantly. On the contrary, P_{aO_2}/F_{IO_2} did not change in O_2 non-responders, while P_{aCO_2} increased in PP from 48 ± 10 mm Hg to 59 ± 15 mm Hg ($P = .01$). Consequently, the VR and pH worsened significantly in PP in this subgroup of subjects (Fig. 2).

Clinical outcomes divided by O_2 responders and non-responders are summarized in Table 3. During the ICU LOS,

Table 1. Population Demographic Characteristics at ICU Admission Divided by Oxygen Response to Pronation

Variables	Total (<i>N</i> = 125)	Oxygen Non-Responders (<i>n</i> = 9)	Oxygen Responders (<i>n</i> = 116)	<i>P</i>
Sex, female	30 (24)	3 (33)	27 (23)	.45
Age, y	62 \pm 11	67 \pm 8	61 \pm 11	.08
Weight, kg	86 \pm 19	74 \pm 9	87 \pm 19	.001
Height, cm	171 \pm 9	169 \pm 7	171 \pm 9	.58
BMI, kg/m ²	30 \pm 6	26 \pm 4	30 \pm 6	.009
Comorbidities				
Hypertension	64 (51)	0	64 (55)	.001
Diabetes	22 (18)	1 (11)	21 (18)	.51
Active smoker	5 (4)	0	5 (4)	.68
Obesity	38 (30)	1 (11)	37 (32)	.18
Cancer	14 (11)	3 (33)	11 (9)	.063
CKD	9 (7)	0	9 (8)	.50
COPD	13 (10)	0	13 (11)	.36
Atrial fibrillation	8 (6)	0	8 (7)	.54
CAD	15 (12)	1 (11)	14 (12)	.71
Liver disease	10 (8)	0	10 (9)	.46
SOFA	5 [4–6]	4 [2–6]	5 [4–7]	.23
SAPS II	38 [33–43]	41 [35–43]	38 [33–44]	.63
Time from onset of symptoms to CT, d	10 \pm 6	14 \pm 7	10 \pm 6	.13
Time from onset of symptoms to first pronation, d	11 \pm 6	15 \pm 6	10 \pm 6	.07

Data are presented as *n* (%), mean \pm SD, or median [interquartile range].

Subjects were defined as oxygen responders to prone position if the P_{aO_2}/F_{IO_2} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. Similarly, oxygen non-responders were defined as those subjects in whom this condition was not satisfied.

BMI = body mass index

CKD = chronic kidney disease

CAD = coronary arterial disease

SOFA = Sequential Organ Failure Assessment

SAPS = Simplified Acute Physiology Score

CT = computed tomography

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

Table 2. Ventilatory Parameters of the Population Divided by Oxygen Response to Pronation in Supine Position Before the First Pronation

Variables	Total (N = 125)	Oxygen Non-Responders (n = 9)	Oxygen Responders (n = 116)	P
ARDS severity				.53
Mild	1 (0.8)	0	1 (1)	
Moderate	65 (52.0)	6 (67)	59 (51)	
Severe	59 (47.2)	3 (33)	56 (48)	
Ventilator setting				
V _T /PBW, mL/kg	6.7 ± 0.9	6.6 ± 1.0	6.8 ± 0.8	.13
Breathing frequency, breaths/min	19 ± 3	20 ± 4	19 ± 3	.86
PEEP, cm H ₂ O	12 ± 2	12 ± 2	12 ± 2	.29
Plateau pressure, cm H ₂ O	24 ± 3	24 ± 3	23 ± 3	.42
C _{RS} , mL/cm H ₂ O	41 ± 11	32 ± 5	42 ± 11	< .001
Driving Pressure, cm H ₂ O	11 ± 3	13 ± 2	11 ± 3	.08
ABG				
pH	7.36 ± 0.07	7.38 ± 0.07	7.36 ± 0.07	.32
P _{aCO₂} , mm Hg	47 ± 9	48 ± 10	47 ± 9	.78
VR	1.6 ± 0.5	1.6 ± 0.4	1.6 ± 0.5	.62
P _{aO₂} , mm Hg	77 ± 16	80 ± 10	77 ± 16	.44
F _{IO₂}	0.8 ± 0.2	0.7 ± 0.2	0.8 ± 0.2	.36
P _{aO₂} /F _{IO₂} , mm Hg	103 [82–123]	113 [89–142]	102 [81–122]	.22

Data are presented as n (%), mean ± SD, or median [interquartile range].

Subjects were defined as oxygen responders to prone position if the P_{aO₂}/F_{IO₂} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. Similarly, oxygen non-responders were defined as those subjects in whom this condition was not satisfied.

V_T = tidal volume

PBW = predicted body weight

C_{RS} = respiratory system compliance

ABG = arterial blood gas

VR = ventilatory ratio

subjects received 4 [2–6] cycles of pronation for a total amount of 80 [46–146] h spent in PP. No differences in ICU LOS ($P = .94$) and survival ($P = .52$) were found between the 2 groups. Bilateral quantitative CT scan analysis did not reveal any difference both in mass and density of hyperinflated, well-aerated, poorly aerated, and non-aerated lung tissue when the subjects were divided into O₂ responders and non-responders (Table 4).

Lastly, the analysis performed on 10 ventral-dorsal lung segments also did not identify any difference between the 2 groups (Fig. 3). For example, the amount of hyperinflated tissue of the ventral region (3.2 ± 2.9 g vs 3.6 ± 2.3 g, $P = .66$) and non-aerated tissue of dorsal regions (254 ± 161 g vs 263 ± 144 g, $P = .85$) were similar between O₂ responders and non-responders. Similar findings were observed when subjects were divided according to the median increase in P_{aO₂}/F_{IO₂} (87 mm Hg). Results can be found in Tables S1, S2, S3, and S4 (see related supplementary materials at <http://www.rcjournal.com>).

CO₂ Response to Pronation

Fifty-one (41%) of 125 subjects improved their VR during PP and were thus defined as CO₂ responders, while the remaining 74 subjects were defined as CO₂ non-responders.

Baseline demographic characteristics are summarized in Table S5 (see related supplementary materials at <http://www.rcjournal.com>). Before PP, CO₂ responders were characterized by higher V_T (6.9 ± 1.0 mL/kg vs 6.6 ± 0.8 mL/kg, $P = .02$), breathing frequency (21 ± 4 breaths/min vs 18 ± 3 breaths/min, $P < .001$), and VR (1.9 ± 0.5 vs 1.5 ± 0.4 , $P < .001$) to maintain a similar arterial P_{CO₂} (49 ± 9 mm Hg vs 46 ± 9 mm Hg, $P = .09$) as compared to CO₂ non-responders (Table S6, see related supplementary materials at <http://www.rcjournal.com>). No differences in days of ventilation ($P = .94$), ICU LOS ($P = .43$), or ICU mortality ($P = .46$) were found between the 2 groups (Table S7, see related supplementary materials at <http://www.rcjournal.com>). Similarly, no differences in quantitative CT results were found between CO₂ responders and non-responders (Table 5).

Discussion

The use of chest CT has permanently changed our understanding of ARDS through its morphological assessment and quantitative analysis of density distribution.¹² For these reasons, this radiological examination is extensively used in some centers to evaluate lung structure, extent of disease, response to lung recruitment,¹⁵ and evolution of disease. In the context of the outbreak of a novel infectious disease

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS




Fig. 2. Variation of clinical parameters during the first pronation cycle of oxygen responders and oxygen non-responders to pronation. Responders were defined as the subjects whose P_{aO_2}/F_{IO_2} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. Panel A represents the variations of respiratory system compliance. Panel B represents the ratio variations between arterial P_{aO_2} and inspiratory fraction of oxygen. Panel C represents the variations in pH. Panel D represents the variations in respiratory ratio. * $P < .050$ vs baseline; \$ $P < .050$ vs non-responders. C_{RS} = respiratory system compliance.

leading to pneumonia and respiratory failure, these concepts were broadly applied. This allowed us to study retrospectively a large number of CT scans of critically ill, mechanically ventilated subjects with COVID-19-related ARDS undergoing PP. Our aim was to evaluate whether the different responses in terms of gas exchange during the first PP were associated with different baseline quantitative CT scan characteristics. In the 125 subjects studied, 93% improved their oxygenation during the first PP, and 41% improved their VR. No relationship was found between quantitative CT scan parameters and either oxygen or CO₂ response to PP. Similar results were observed when dividing the population according to the median increase in P_{aO_2}/F_{IO_2} . Moreover, we confirmed that the O₂ responders were characterized by higher baseline compliance and lower driving pressure as compared to non-responders. Of note, the time spent in PP did not differ between the 2 groups.

During the COVID-19 pandemic, pronation was broadly used in mechanically ventilated patients.^{1,29-31} In line with

previous data,¹ most of the studied subjects improved their oxygenation during the first PP. Aalinezhad and colleagues³² identified in subjects with COVID-19 a relationship between the severity of lung involvement measured at CT scan and blood oxygenation.³² Moreover, the possibility of predicting lung recruitment from a single static baseline CT scan using a machine learning approach has recently been described.³³ In classic ARDS, lung perfusion is similar in prone and supine position, being slightly unbalanced toward dorsal lung regions.^{34,35} According to this characteristic, the improvement in P_{aO_2}/F_{IO_2} in PP should parallel the variation in density distribution, corresponding to an increase in well-aerated lung tissue in the dorsal areas of the lungs. Despite these premises, in classic ARDS, Papazian and colleagues³⁶ found no correlation between baseline quantitative CT data and P_{aO_2}/F_{IO_2} response to pronation. In addition, similarly to what has been observed in non-intubated subjects with COVID-19,¹⁸ we were not able to identify a correlation between baseline quantitative CT-scan characteristics and P_{aO_2}/F_{IO_2} response during PP.

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

Table 3. Clinical Outcomes of the Population Divided by Oxygen Response to Pronation

Variables	Total (N = 125)	Oxygen Non-Responders (n = 9)	Oxygen Responders (n = 116)	P
Pronation sessions, no.	4 [2–6]	3 [2–5]	4 [2–7]	.27
Total pronation time, h	80 [46–146]	92 [47–139]	79 [44–148]	.82
INO	32 (26)	3 (33)	29 (25)	.69
Duration of ventilation, d	30 [17–42]	42 [19–57]	29 [17–39]	.25
Tracheostomy	76 (61)	5 (56)	71 (61)	.74
Hospital LOS, d	45 [26–65]	57 [22–62]	44 [26–65]	.94
ICU LOS, d	33 [19–45]	43 [10–50]	33 [20–45]	.94
ICU Outcome				.52
Deceased	51 (41)	4 (44)	47 (40)	
Discharged	74 (59)	5 (56)	70 (60)	

Data are presented as *n* (%) or median [interquartile range].

Subjects were defined as oxygen responders to prone position if the P_{aO_2}/F_{IO_2} ratio increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. Similarly, oxygen non-responders were defined as those subjects in whom this condition was not satisfied.

INO = inhaled nitric oxide

LOS = length of stay

Table 4. Baseline Quantitative Computed Tomography Parameters of the Population Divided for the Oxygen Response to Pronation

Variables	Total (N = 125)	Oxygen Non-Responders (n = 9)	Oxygen Responders (n = 116)	P
Bilateral lung				
Volume, mL	3,526 \pm 1,009	3,816 \pm 757	3,503 \pm 1,025	.27
Density, HU	–545 \pm 106	–562 \pm 58	–544 \pm 109	.42
Tissue mass, g	1,541 \pm 390	1,654 \pm 304	1,532 \pm 396	.28
Hyperinflated lung				
Volume, mL	160 \pm 157	189 \pm 126	158 \pm 159	.50
Density, HU	–968 \pm 2	–968 \pm 2	–968 \pm 2	.72
Tissue mass, g	5 \pm 5	6 \pm 3	5 \pm 5	.46
Well-aerated lung				
Volume, mL	2,092 \pm 895	2,283 \pm 532	2,077 \pm 917	.31
Density, HU	–740 \pm 31	–752 \pm 17	–739 \pm 32	.066
Tissue mass, g	530 \pm 208	568 \pm 13	527 \pm 213	.43
Poorly aerated lung				
Volume, mL	891 \pm 324	939 \pm 242	887 \pm 330	.56
Density, HU	–343 \pm 20	–345 \pm 16	–343 \pm 20	.75
Tissue mass, g	584 \pm 210	615 \pm 158	582 \pm 214	.57
Non-aerated lung				
Volume, mL	319 \pm 192	335 \pm 157	317 \pm 195	.75
Density, HU	–51 \pm 8	–50 \pm 6	–51 \pm 8	.75
Tissue mass, g	303 \pm 185	319 \pm 150	302 \pm 188	.76

Data are presented as mean \pm SD.

Subjects were defined as oxygen responders to prone position if the P_{aO_2}/F_{IO_2} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. Similarly, oxygen non-responders were defined as those subjects in whom this condition was not satisfied.

The negative findings of these studies might have several explanations. CT data accurately describe lung parenchymal density, while they do not assess pulmonary perfusion. This aspect might be of utmost importance in patients with COVID-19–associated ARDS. Indeed, this disease is characterized by (1) impairment of hypoxic vasoconstriction

leading to a marked ventilation/perfusion mismatch^{37–39} and (2) the diffuse presence of pulmonary microthrombosis.⁴⁰ Since both of these vascular defects can be diffused to all the lungs, irrespective of gravitational distribution (dependent vs non-dependent), and regardless of the parenchymal aspect assessed with CT scan, a dissociation between

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS




Fig. 3. Ventral-dorsal (1 to 10 segment) regional lung tissue distribution subjects divided by oxygen responders and non-responders to pronation. O₂ responders defined as the subjects whose P_{aO₂}/F_{IO₂} increased by ≥ 20 mm Hg during prone ventilation as compared to baseline values in supine position. A: Hyperinflated tissue, B: well aerated tissue, C: poorly aerated tissue, and D: non-aerated tissue.

Table 5. Baseline Quantitative Computed Tomography Parameters of the Population Divided for the CO₂ Response to Pronation

Variables	Total (N = 125)	CO ₂ Non-Responders (n = 74)	CO ₂ Responders (n = 51)	P
Bilateral lung				
Volume, mL	3,526 ± 1,009	3,502 ± 1,030	3,560 ± 987	.75
Density, HU	-545 ± 106	-546 ± 94	-545 ± 123	.99
Tissue mass, g	1,541 ± 390	1,534 ± 377	1,551 ± 412	.89
Hyperinflated lung				
Volume, mL	160 ± 157	144 ± 136	184 ± 182	.37
Density, HU	-968 ± 2	-968 ± 2	-968 ± 2	.58
Tissue mass, g	5 ± 5	4 ± 4	6 ± 5	.36
Well-aerated lung				
Volume, mL	2,092 ± 895	2,087 ± 885	2,099 ± 918	.96
Density, HU	-740 ± 31	-735 ± 29	-748 ± 33	.02
Tissue mass, g	530 ± 208	542 ± 216	512 ± 198	.55
Poorly aerated lung				
Volume, mL	891 ± 324	929 ± 323	835 ± 320	.07
Density, HU	-343 ± 20	-346 ± 19	-338 ± 21	.033
Tissue mass, g	584 ± 210	606 ± 207	553 ± 212	.10
Non-aerated lung				
Volume, mL	319 ± 192	287 ± 157	366 ± 228	.13
Density, HU	-51 ± 8	-52 ± 6	-49 ± 9	.19
Tissue mass, g	304 ± 185	272 ± 150	349 ± 220	.12

Data are presented as mean ± SD.

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

aeration/ventilation and gas exchange has been described in subjects with COVID-19–associated ARDS.³⁸ It is thus conceivable that, in addition to the unknown potential for lung recruitment, the variable and unpredictable lung perfusion changes further hinder the prediction of the response solely based on baseline quantitative CT information. In addition, the varying potential involvement of pulmonary vasculature could justify the broad spectrum of oxygen response to pronation reported, ranging from 35% to 93%.^{1,2,13,18,24,41} Interestingly, and in line with this reasoning, in our population no difference in quantitative CT characteristics was observed between subjects with moderate vs severe COVID-19–associated ARDS (Table S8, see related supplementary materials at <http://www.rcjournal.com>).

A second possible explanation of the different responses to pronation might be the disease time course. Indeed, despite the lack of statistical significance, the time between symptom onset and first pronation was longer in O₂ non-responders, possibly resulting in a more severe disease stage, as suggested by the lower C_{RS}. In line with this hypothesis, a decreasing response to PP (in terms of oxygenation) has been described as a consequence of lung consolidation toward organizing fibrotic pneumonia.^{13,42,43} Regardless of this potential explanation, no relevant differences in gas exchange, lung weight, or non-aerated lung tissue were noted between O₂ responders and non-responders.

Response to Pronation

In response to PP, O₂ responders increased slightly, but significantly their C_{RS}, which remained above baseline after re-supination. The improvement of P_{aO₂}/F_{IO₂} paralleled C_{RS}, except for a slight decrease after re-supination. Notably, these variations were not mirrored by the VR, which did not change significantly.

Fossali et al⁴⁴ performed a physiological study exploring the early changes after pronation in subjects with COVID-19–associated ARDS, performing CT scans both in supine and PPs. They demonstrated that PP significantly decreased the weight of non-aerated and hyperinflated lung tissue and increased the amount of normally aerated lung. Moreover, the regional response to PP was not homogenous, as demonstrated by the remarkable recruitment in the dorsal regions and de-recruitment in the ventral. However, in our population, O₂ responders and non-responders, despite having similar baseline amounts of hyperinflated ventral tissue and non-aerated dorsal tissue, demonstrated markedly different responses in terms of lung mechanics and gas exchanges during PP. Notably, also in Fossali's⁴⁴ work, no association between the amount of ventral de-recruitment or dorsal recruitment and the oxygen response was found. Taken together, our results and the findings of this author foster two considerations: *First*, in patients with COVID-19, the oxygen response to pronation is most likely not

predictable from a static baseline CT scan; and *second*, the observed increase in oxygenation is possibly due mainly to the improvement of the ventilation/perfusion matching related to a persistency of perfusion in the vertebral part of the thorax and a reopening of the dorsal collapsed lung.⁴⁵⁻⁴⁸ This second hypothesis is corroborated by the study of Richter et al,⁴⁶ who demonstrated that the oxygenation response to pronation in subjects with ARDS was consequent to an improvement in ventilation/perfusion match due to the unchanged perfusion in the dorsal part of the thorax associated with a reopening of the dorsal collapsed lung. This mechanism, also described using the ventilation/perfusion tools of electrical impedance tomography in classic subjects with ARDS,⁴⁹ was confirmed in COVID-19–associated ARDS.^{50,51}

In O₂ non-responders, no variations in C_{RS} or P_{aO₂}/F_{IO₂} were observed. Furthermore, differently from responders, they experienced an increase in VR and P_{aCO₂} that persisted after the re-supination. This observation points toward an increased dead space. As an increase in hyperinflation is unlikely the underlying mechanism, we think that also in this case the worsening of ventilation/perfusion might be the cause. Of note, also when dividing the population according to their response in terms of CO₂ clearance (ie, variation in VR) no difference in baseline quantitative CT data was observed.

Automated CT Scan Segmentation

Despite the absence of association between CT scan characteristics and response to PP, CT exams represent the accepted standard for evaluating the alterations of lung parenchyma, even in the early stages of disease, when the patient has few or no symptoms. Moreover, it is also a useful tool for monitoring disease progression.^{11,52}

In addition to the classic qualitative visual image interpretation, the automated and integrated workflow of image analysis allows to extract several objective metadata quantitative information retained in the image, such as parenchymal density and volume, and permits the definition of lung compartments based on the different degrees of aeration.¹² Although CT image analysis could be extremely informative, some aspects need to be considered for its use in clinical practice. In this work, an algorithm for automatic segmentation of lungs in CT images was employed,²⁵ which drastically reduces analysis time and enables real-time quantitative results through the use of dedicated in-house software. Avoiding the time-consuming task of drawing the lung boundaries (selection of regions of interest), the physician can thus focus more on interpreting the results of the obtained quantitative metrics.

Limitations

Several limitations have to be addressed for this study. *First*, due to the retrospective nature of the study and the low number of O₂ non-responders, our analyses could be

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

underpowered to identify any difference in quantitative CT characteristics; a controlled methodology and more homogeneous groups may produce different results. *Second*, the CT scans were performed for clinical purposes and retrospectively used for the analyses. Thus, no standardization of ventilation mode nor respiratory phase (eg, inspiratory pause) was performed during CT scan acquisition. *Third*, no data regarding perfusion of the lung were available. Consequently, the pathophysiological role of ventilation/perfusion matching in explaining oxygen and CO₂ responses to pronation can only be hypothesized. *Finally*, the trunk inclination angle used during the respiratory mechanics measurement was not standardized.⁵³⁻⁵⁵

Conclusions

Most subjects with COVID-19–related ARDS improved their oxygenation at the first pronation cycle. Our study performed on a large population of critically ill, mechanically ventilated subjects with COVID-19–related ARDS suggests that quantitative data obtained from a baseline CT scan are neither associated with the oxygen response nor with the response in terms of CO₂ elimination.

ACKNOWLEDGMENTS

The authors are deeply grateful to all the physicians and nurses of the COVID-19 ICUs of the Niguarda Hospital.

REFERENCES

- Langer T, Brioni M, Guzzardella A, Carlesso E, Cabrini L, Castelli G, et al; PRONA-COVID Group. Prone position in intubated, mechanically ventilated patients with COVID-19: a multi-centric study of more than 1,000 patients. *Crit Care* 2021;25(1):1-11.
- Protti A, Santini A, Pennati F, Chiurazzi C, Ferrari M, Iapichino GE, et al. Lung response to prone positioning in mechanically-ventilated patients with COVID-19. *Crit Care* 2022;26(1):127-129.
- Gattinoni L, Taccone P, Carlesso E, Marini JJ. Prone position in acute respiratory distress syndrome. Rationale, indications, and limits. *Am J Respir Crit Care Med* 2013;188(11):1286-1293.
- Le MQ, Rosales R, Shapiro LT, Huang LY. The down side of prone positioning: the case of a coronavirus 2019 survivor. *Am J Phys Med Rehabil* 2020;99(10):870-872.
- Zadek F, Rubin J, Grassi L, Van Den Kroonenberg D, Larson G, Capriles M, et al. Individualized multimodal physiologic approach to mechanical ventilation in patients with obesity and severe acute respiratory distress syndrome reduced venovenous extracorporeal membrane oxygenation utilization. *Crit Care Explor* 2021;3(7):e0461.
- Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving sepsis campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Intensive Care Med* 2020;46(5):854-887.
- Lassola S. Pronation in acute respiratory distress syndrome (ARDS) secondary to COVID-19. *J Public Heal Emerg* 2022;6(1):1-9.
- Fang Y, Zhang H, Xie J, Lin M, Ying L, Pang P, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology* 2020;296(2):E115-E117.
- Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, et al. Correlation of chest CT and RT-PCR testing for coronavirus disease 2019 (COVID-19) in China: a report of 1,014 cases. *Radiology* 2020;296(2):E32-E40.
- Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* 2020;20(4):425-434.
- Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 2020;295(1):202-207.
- Gattinoni L, Caironi P, Pelosi P, Goodman LR. What has computed tomography taught us about the acute respiratory distress syndrome? *Am J Respir Crit Care Med* 2001;164(9):1701-1711.
- Rossi S, Palumbo MM, Sverzellati N, Busana M, Malchiodi L, Bresciani P, et al. Mechanisms of oxygenation responses to proning and recruitment in COVID-19 pneumonia. *Intensive Care Med* 2022;48(1):56-66.
- Henzler D, Mahnken AH, Wildberger JE, Rossaint R, Günther RW, Kühlen R. Multi-slice spiral computed tomography to determine the effects of a recruitment maneuver in experimental lung injury. *Eur Radiol* 2006;16(6):1351-1359.
- Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri VM, Quintel M, et al. Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med* 2006;354(17):1775-1786.
- Bugedo G, Bruhn A, Hernández G, Rojas G, Varela C, Tapia JC, et al. Lung computed tomography during a lung recruitment maneuver in patients with acute lung injury. *Intensive Care Med* 2003;29(2):218-225.
- Chen YY, Kuo JSH, Ruan SY, Chien YC, Ku SC, Yu CJ, et al. Prognostic value of computed tomographic findings in acute respiratory distress syndrome and the response to prone positioning. *BMC Pulm Med* 2022;22(1):71-12.
- Raimondi F, Cazzaniga S, Annibaldi S, Novelli L, Brivio M, Pappacena S, et al. Extent and distribution of parenchymal abnormalities in baseline CT scans do not predict awake prone positioning response in COVID-19–related ARDS. *Diagnostics (Basel)* 2022;12(8):1848.
- Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin definition. *JAMA* 2012;307(23):2526-2533.
- Foti G, Giannini A, Bottino N, Castelli Gian P, Cecconi M, Grasselli G, et al; COVID-19 Lombardy ICU Network. Management of critically ill patients with COVID-19: suggestions and instructions from the coordination of intensive care units of Lombardy. *Minerva Anestesiol* 2020;86(11):1234-1245.
- Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al; COVID-19 Lombardy ICU Network. Risk factors associated with mortality among patients with COVID-19 in intensive care units in Lombardy, Italy supplemental content. *JAMA Intern Med* 2020;180(10):1345-1355.
- Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A; Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342(18):1301-1308.
- Sinha P, Fauvel NJ, Singh S, Soni N. Ventilatory ratio: a simple bedside measure of ventilation. *Br J Anaesth* 2009;102(5):692-697.
- Scaramuzza G, Gamberini L, Tonetti T, Zani G, Ottaviani I, Mazzoli CA, et al; ICU-RER COVID-19 Collaboration. Sustained oxygenation improvement after first prone positioning is associated with liberation from mechanical ventilation and mortality in critically ill COVID-19 patients: a cohort study. *Ann Intensive Care* 2021;11(1):63-10.
- Berta L, Rizzetto F, De Mattia C, Lizio D, Felisi M, Colombo PE, et al; Niguarda COVID-19 Working Group. Automatic lung

QUANTITATIVE CT AND PRONATION IN COVID-19 ARDS

- segmentation in COVID-19 patients: impact on quantitative computed tomography analysis. *Phys Med* 2021;87(6):115-122.
26. Berta L, De Mattia C, Rizzetto F, Carrazza S, Colombo PE, Fumagalli R, et al. A patient-specific approach for quantitative and automatic analysis of computed tomography images in lung disease: application to COVID-19 patients. *Phys Med* 2021;82(12):28-39.
 27. Zorzi G, Berta L, Rizzetto F, De Mattia C, Felisi MMJ, Carrazza S, et al. Artificial intelligence for differentiating COVID-19 from other viral pneumonias on CT: comparative analysis of different models based on quantitative and radiomic approaches. *Eur Radiol Exp* 2023;7(1):3.
 28. Vecchi V, Langer T, Bellomi M, Rampinelli C, Chung KK, Cancio LC, et al. Low-dose CT for quantitative analysis in acute respiratory distress syndrome. *Crit Care* 2013;17(4):R183.
 29. Guerin C, Baboi L, Richard JC. Mechanisms of the effects of prone positioning in acute respiratory distress syndrome. *Intensive Care Med* 2014;40(11):1634-1642.
 30. Mentzelopoulos SD, Roussos C, Zakynthinos SG. Prone position reduces lung stress and strain in severe acute respiratory distress syndrome. *Eur Respir J* 2005;25(3):534-544.
 31. Albert RK, Keniston A, Baboi L, Ayzac L, Guérin C; PROSEVA Investigators. Prone position-induced improvement in gas exchange does not predict improved survival in the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2014;189(4):494-496.
 32. Aalinezhad M, Alikhani F, Akbari P, Rezaei MH, Soleimani S, Hakamifard A. The relationship between CT severity score and capillary blood oxygen saturation in patients with COVID-19 infection. *Indian J Crit Care Med* 2021;25(3):279-283.
 33. Pennati F, Aliverti A, Pozzi T, Gattarello S, Lombardo F, Coppola S, et al. Machine learning predicts lung recruitment in acute respiratory distress syndrome using single lung CT scan. *Ann Intensive Care* 2023;13(1):60.
 34. Musch G, Layfield JDH, Harris RS, Vidal Melo MF, Winkler T, Callahan RJ, et al. Topographical distribution of pulmonary perfusion and ventilation, assessed by PET in supine and prone humans. *J Appl Physiol* (1985) 2002;93(5):1841-1851.
 35. Henderson AC, Sá RC, Theilmann RJ, Buxton RB, Prisk GK, Hopkins SR. The gravitational distribution of ventilation/perfusion ratio is more uniform in prone than supine posture in the normal human lung. *J Appl Physiol* (1985) 2013;115(3):313-324.
 36. Papazian L, Paladini MH, Bregeon F, Thirion X, Durieux O, Gannier M, et al. Can the tomographic aspect characteristics of patients presenting with acute respiratory distress syndrome predict improvement in oxygenation-related response to the prone position? *Anesthesiology* 2002;97(3):599-607.
 37. Si-Mohamed S, Chebib N, Sigovan M, Zumbihl L, Turquier S, Bocalini S, et al. In vivo demonstration of pulmonary microvascular involvement in COVID-19 using dual-energy computed tomography. *Eur Respir J* 2020;56(4):2002608.
 38. Santamarina MG, Boisier D, Contreras R, Baque M, Volpacchio M, Beddings I. COVID-19: a hypothesis regarding the ventilation/perfusion mismatch. *Crit Care* 2020;24(1):395-397.
 39. Busana M, Giosa L, Cressoni M, Gasperetti A, Di Girolamo L, Martinelli A, et al. The impact of ventilation/perfusion inequality in COVID-19: a computational model. *J Appl Physiol* (1985) 2021;130(3):865-876.
 40. Patel BV, Arachchillage DJ, Ridge CA, Bianchi P, Doyle JF, Garfield B, et al. Pulmonary angiopathy in severe COVID-19: physiologic, imaging, and hematologic observations. *Am J Respir Crit Care Med* 2020;202(5):690-699.
 41. Sanabria-Rodríguez OO, Cardozo-Avendaño SL, Muñoz-Velandia OM. Factors associated with a nonresponse to prone positioning in patients with severe acute respiratory distress syndrome due to SARS-CoV-2. *Crit Care Sci* 2023;35(2):156-162.
 42. Barisione E, Grillo F, Ball L, Bianchi R, Grosso M, Morbini P, et al. Fibrotic progression and radiologic correlation in matched lung samples from COVID-19 post-mortems. *Virchows Arch* 2021;478(3):471-485.
 43. Patel BV, Haar S, Handlip R, Auepanwiriyaikul C, Lee TML, Patel S, et al; United Kingdom COVID-ICU National Service Evaluation. Natural history, trajectory, and management of mechanically ventilated COVID-19 patients in the United Kingdom. *Intensive Care Med* 2021;47(5):549-565.
 44. Fossali T, Pavlovsky B, Ottolina D, Colombo R, Basile MC, Castelli A, et al. Effects of prone position on lung recruitment and ventilation/perfusion matching in patients with COVID-19 acute respiratory distress syndrome: a combined CT scan/electrical impedance tomography study. *Crit Care Med* 2022;50(5):723-732.
 45. Guérin C, Reignier J, Richard J-C, Beuret P, Gacouin A, Boulain T, et al; PROSEVA Study Group. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368(23):2159-2168.
 46. Richter T, Bellani G, Harris RS, Melo MFV, Winkler T, Venegas JG, et al. Effect of prone position on regional shunt, aeration, and perfusion in experimental acute lung injury. *Am J Respir Crit Care Med* 2005;172(4):480-487.
 47. Lee DL, Chiang HT, Lin SL, Ger LP, Kun MH, Huang YCT. Prone-position ventilation induces sustained improvement in oxygenation in patients with acute respiratory distress syndrome who have a large shunt. *Crit Care Med* 2002;30(7):1446-1452.
 48. Lamm WJE, Graham MM, Albert RK. Mechanism by which the prone position improves oxygenation in acute lung injury. *Am J Respir Crit Care Med* 1994;150(1):184-193.
 49. Wang Y, Zhong M, Dong M, Song J, Zheng Y, Wu W, et al. Prone positioning improves ventilation/perfusion matching assessed by electrical impedance tomography in patients with ARDS: a prospective physiological study. *Crit Care* 2022;26(1):154.
 50. Zarantonello F, Andreatta G, Sella N, Navalesi P. Prone position and lung ventilation and perfusion matching in acute respiratory failure due to COVID-19. *Am J Respir Crit Care Med* 2020;202(2):278-279.
 51. Perier F, Tuffet S, Maraffi T, Alcalá G, Victor M, Haudebourg AF, et al. Effect of positive end-expiratory pressure and proning on ventilation and perfusion in COVID-19 acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020;202(12):1713-1717.
 52. Rubin GD, Ryerson CJ, Haramati LB, Sverzellati N, Kanne JP, Raouf S, et al. The role of chest imaging in patient management during the COVID-19 pandemic: a multinational consensus statement from the Fleischner Society. *Chest* 2020;158(1):106-116.
 53. Marrazzo F, Spina S, Zadek F, Forlini C, Bassi G, Giudici R, et al. PEEP titration is markedly affected by trunk inclination in mechanically ventilated patients with COVID-19 ARDS: a physiologic, crossover study. *J Clin Med* 2023;12(12):3914.
 54. Marrazzo F, Spina S, Forlini C, Guarnieri M, Giudici R, Bassi G, et al. Effects of trunk inclination on respiratory mechanics in patients with COVID-19-associated acute respiratory distress syndrome: let's always report the Angle! *Am J Respir Crit Care Med* 2022;205(5):582-584.
 55. Marrazzo F, Spina S, Zadek F, Forlini C, Bassi G, Giudici R, et al. Ventilation distribution during changes in trunk inclination in patients with ARDS. *Respir Care* 2024;69(2):222-226.