

Alveolar Recruitment Maneuvers Under General Anesthesia: A Systematic Review of the Literature

Benjamin L Hartland RN, Timothy J Newell RN, and Nicole Damico MSNA CRNA

Introduction

- Description of Condition**
- Description of Interventions**
- Significance of Review**

Objectives

Methods

- Types of Studies**
- Types of Subjects**
- Search Methods for Identification of Studies**

Data Collection and Analysis

- Selection of Studies**
- Assessment of Risk of Bias in Included Studies**
- Sensitivity Analysis**

Results

- Results of Search**
- Included Studies**
- Risk of Bias of Included Studies**
- Primary Outcomes**
- Secondary Outcomes**

Discussion

- Summary of Main Results**
- Primary Outcome Measures**
- Secondary Outcome Measures**
- Major Clinical Themes**
- Overall Completeness and Applicability of Evidence**
- Quality of Evidence**
- Potential Bias in the Review Process**

Conclusions

- Implications for Practice**
- Implications for Research**

BACKGROUND: The sigh is a normal homeostatic reflex that maintains lung compliance and decreases atelectasis. General anesthesia abolishes the sigh reflex with rapid onset of atelectasis in 100% of patients. Studies show a strong correlation between atelectasis and postoperative pulmonary complications, raising health-care costs. Alveolar recruitment maneuvers recruit collapsed alveoli, increase gas exchange, and improve arterial oxygenation. There is no consensus in the literature about the benefits of alveolar recruitment maneuvers. A systematic review is necessary to delineate their usefulness. **METHODS:** The search strategy included utilizing PubMed, CINAHL, the Cochrane Library, the National Guideline Clearinghouse, and all subsequent research reference lists up to January 2014. Inclusion criteria involved studies that compared the use of an alveolar

recruitment maneuver with a control group lacking an alveolar recruitment maneuver in adult surgical subjects not suffering from ARDS or undergoing cardiac or thoracic surgeries. **RESULTS:** Six randomized controlled trials of the 439 studies initially identified achieved a score of ≥ 3 on the Jadad scale and were included in this review. Alveolar recruitment maneuvers consisted of a stepwise increase in tidal volume to a plateau pressure of 30 cm H₂O, a stepwise increase in PEEP to 20 cm H₂O, or sustained manual inflations of the anesthesia reservoir bag to a peak inspiratory pressure of 40 cm H₂O. Subjects in the alveolar recruitment maneuver groups experienced a higher intraoperative P_{aO₂} with improved lung compliance. Different alveolar recruitment maneuvers were equally effective. There was a significant advantage when alveolar recruitment maneuvers were followed by PEEP application. **CONCLUSIONS:** Alveolar recruitment maneuvers followed by PEEP should be instituted after induction of general anesthesia, routinely during maintenance, and in the presence of a falling S_{pO₂} whenever feasible. They allow the anesthesia provider to reduce the F_{IO₂} while maintaining a higher S_{pO₂}, limiting the masking of shunts. Utilization of alveolar recruitment maneuvers may reduce postoperative pulmonary complications and improve patient outcomes. *Key words:* sigh breaths; alveolar recruitment maneuvers; lung-protective ventilation. [Respir Care 2015;60(3):1-9. © 2015 Daedalus Enterprises]

Introduction

Description of Condition

In 1964, Bendixen et al¹ found that awake adults sigh an average of 9–10 times/h. The sigh breath is a normal homeostatic reflex characterized by a complex interaction between vagally mediated input and peripheral chemoreceptors that results in an increase in inspiration.² The sigh breath functions to maintain pulmonary compliance.³ In a group of spontaneously breathing subjects, pulmonary compliance decreased in the absence of periodic sighs and returned to baseline after only 2 sigh breaths. Sigh breaths also minimize the alveolar-arterial oxygen difference (P_{(A-a)O₂}) and maintain venous admixture within normal range.⁴

Sigh breaths help to release new surfactant and distribute it evenly on the alveolar surface and the distal airways.^{5,6} The fusion pore is the area on the alveolar type II cell that acts as a mechanical barrier to prevent further surfactant release. Sigh breaths open this barrier and replenish the available supply of surfactant.⁶

The authors are affiliated with the Department of Nurse Anesthesia, Virginia Commonwealth University, Richmond, Virginia.

Supplementary material related to this paper is available at <http://www.rcjournal.com>.

The authors have disclosed no conflicts of interest.

Correspondence: Benjamin L Hartland RN-BSN, Department of Nurse Anesthesia, Virginia Commonwealth University, 1200 E Broad Street, Richmond, VA 23298. E-mail: hartlandbl@vcu.edu.

DOI: 10.4187/respcare.03488

If sigh breaths occur in everyday life to prevent atelectasis as a normal homeostatic reflex, what happens when patients receive neuromuscular blocking drugs, are intubated, and are exposed to high concentrations of oxygen with a fixed tidal volume (V_T) on a ventilator? General endotracheal anesthesia causes compression and absorption atelectasis through 3 main mechanisms: dyskinesia, increased F_{IO₂}, and ablation of the sigh breath. Dyskinesia causes a restriction in the movement of a patient's dependent diaphragm, resulting in a decrease in lung compliance.⁷ It also decreases movement of the dependent portion of the lungs, thereby decreasing functional residual capacity.⁸ These factors contribute to compression atelectasis, which occurs when the pressure distending the alveolus is reduced.⁹ In a study using computed tomography, atelectasis was noted to occur in 100% of subjects undergoing general anesthesia.¹⁰

Absorption atelectasis occurs when the flux of oxygen from alveolar gas into capillary blood exceeds waste gas returning to the alveoli (as oxygen diffuses faster than nitrogen). This frequently occurs during induction of anesthesia when the F_{IO₂} is increased to 100%. Since oxygenated gas leaves the alveoli for the blood faster than waste gas (mostly nitrogen) returns to the alveoli, the alveoli shrink and eventually collapse, increasing atelectasis.¹¹

In 1964, Bendixen et al⁴ hypothesized that constant ventilation with adequate but static V_T in anesthetized patients would result in progressive atelectasis and shunting when sigh breaths were absent. They found that, on average, P_{aO₂} fell 22% and pulmonary compliance fell 15% in the absence of sigh breaths. After a few minutes of slow, deep, sustained breaths, P_{aO₂} rose an average of 150 mm Hg, reducing the shunt created by static V_T. The law of Laplace explains the increase in P_{aO₂}: pressure = 2T/r, where

T = surface tension, and r = radius. When the radius of an alveolus is decreased, the pressure required to re-inflate that alveolus increases. Alveolar recruitment maneuvers provide the elevated pressure necessary to re-inflate collapsed alveoli and may help to prevent impending collapse.

Induction and maintenance of anesthesia with muscle paralysis and a fixed V_T eliminate a patient's ability to generate the sigh breath. Without this inherent ability to decrease atelectasis and improve gas exchange, the patient is vulnerable to the consequences of atelectasis.

Description of Interventions

Alveolar recruitment maneuvers are described as vital capacity breaths, double V_T breaths, and sigh breaths.¹² Sigh breaths are to awake, spontaneously breathing patients as alveolar recruitment maneuvers are to anesthetized, mechanically ventilated patients. They use sustained increases in airway pressure (breath-holds) to recruit collapsed alveoli and improve arterial oxygenation.¹³

The method for achieving an alveolar recruitment maneuver tends to fall into 2 groups. Some authors consider sustained inflation of the lungs for 5–30 s to a prescribed peak inspiratory pressure (PIP) as representative of alveolar recruitment maneuvers.^{14–18} Others describe incrementally increasing PEEP in a stepwise manner as an alveolar recruitment maneuver.^{19,20} Both methods include the use of PEEP following an alveolar recruitment maneuver to maintain its benefit. For the purpose of this review, studies utilizing both methods will be used to determine the most effective application of an alveolar recruitment maneuver.

Significance of Review

Atelectasis likely occurs in up to 100% of patients undergoing general anesthesia.¹⁰ The intraoperative effects of atelectasis include an increase in the $P_{(A-a)O_2}$, an increase in pulmonary shunting, and a decrease in S_{pO_2} .^{4,15} Atelectasis may increase the risk of volutrauma as a constant volume is imparted to a dwindling alveolar volume. Postoperative pulmonary complications are defined as any respiratory condition that adversely influences the clinical course of patients after surgery.²¹ Studies show a strong correlation between atelectasis and postoperative pulmonary complications. Atelectasis appears to be one of the primary mechanisms underlying acute lung injury, is a major cause of postoperative hypoxemia, and is associated with a prolonged ICU and hospital stay.²¹ Atelectasis may also contribute to serious postoperative pulmonary complications such as respiratory failure and pneumonia.²² Following cardiac bypass surgery, atelectasis causes hypoxemia and pulmonary shunting.²³ Postoperative pulmonary complications represent a substantial economic burden due

to longer hospital stays and increased health-care costs.²¹ In 2010, postoperative pulmonary complications were estimated to add nearly 3.5 billion dollars to annual health-care costs.²⁴

Despite their proposed benefits, there is no general consensus in the literature regarding the appropriate methods, timing, evaluation, and benefits of alveolar recruitment maneuvers during general anesthesia. Study outcomes range from improving arterial oxygenation, to no effect, to decreases in cardiac output and blood pressure.^{19,25} A thorough and focused systematic review regarding alveolar recruitment maneuvers is necessary to delineate their usefulness in improving patient outcomes both during and after general anesthesia.

Objectives

The primary objectives of this systematic review were: (1) to identify existing evidence in the literature; (2) to determine the usefulness, appropriate method, surgical population, and timing of alveolar recruitment maneuvers in the intraoperative period; (3) to determine the role of PEEP in alveolar recruitment maneuvers; (4) to identify methods for evaluating the efficacy of alveolar recruitment maneuvers; and (5) to highlight areas of future research related to alveolar recruitment maneuvers.

Methods

Types of Studies

Studies published before January 2014 were reviewed and were not limited by language. Studies based on subjects with ARDS were excluded because these subjects often require the use of different ventilatory strategies that may limit or skew the effects of standard alveolar recruitment maneuvers. Studies involving cardiac and thoracic surgeries were excluded because of surgical factors impacting normal heart and lung physiology. Animal studies were excluded because of the questionable generalizability of findings. To limit the effects of bias and covariates, only systematic reviews, quantitative meta-analyses, and randomized controlled trials (RCTs) were included. Finally, studies were limited to the intraoperative period, with assessment continuing in the immediate postoperative period for subjects undergoing general endotracheal anesthesia (Fig. 1).

Types of Subjects

Only adult subjects age 16 y and older were represented in these studies because of potential differences stemming from neonatal and pediatric anatomies. Alveolar recruitment maneuvers included either a stepwise increase in




Fig. 1. Flow chart. RCT = randomized controlled trial. ETGA = endotracheal general anesthesia.

Table 1. Primary and Secondary Outcome Measures

Primary Outcome Measures	Secondary Outcome Measures
Intraoperative P_{aO_2}	Postoperative pulmonary complications
Pulmonary compliance	Postoperative P_{aO_2} or S_{pO_2}
$P_{(A-a)O_2}$	Alveolar recruitment maneuver complications
P_{aO_2}/F_{IO_2}	Alveolar recruitment maneuver frequency
	Airway resistance

$P_{(A-a)O_2}$ = alveolar-arterial oxygen difference

PEEP or V_T using a ventilator or manual inflation of a subject's lungs to a predetermined PIP using an anesthesia reservoir bag. Each treatment group was compared with a control group that did not receive an alveolar recruitment maneuver. See Table 1 for a description of outcome measures.

Search Methods for Identification of Studies

The search included PubMed, CINAHL, the Cochrane Library, and the National Guideline Clearinghouse. Additional studies were detected by searching the reference

lists of all included research reports. Search terms included: lung recruitment measures, alveolar recruitment measures, perioperative recruitment maneuvers, PEEP, general anesthesia and chest x-rays, preventing atelectasis during anesthesia/anesthesia, atelectasis and shunting during anesthesia/anesthesia, lung atelectasis and anesthesia/anesthesia, anesthesia/anesthesia and atelectasis development, sigh breaths, sigh breaths during ventilation, lung recruitment maneuvers, and perioperative atelectasis.

Data Collection and Analysis

Selection of Studies

Two reviewers (BLH and TJN) independently assessed all potential studies from the search strategy for inclusion. Consensus was reached through discussion or, if required, by consultation with a third reviewer (ND). The reviewers were not blinded to the authors or results of the studies. Data were independently extracted by the reviewers and cross-checked. Consensus was reached through discussion or, if required, by consultation with the third reviewer. A standard data extraction form was used (see the supplementary materials at <http://www.rcjournal.com>).

Assessment of Risk of Bias in Included Studies

The Jadad scale was used to evaluate methodological quality.²⁶ The methodological quality assessment was performed by 2 reviewers as stated above. In cases of a discrepancy in the scores assigned by the primary reviewers, the third reviewer was used to reach a consensus. The Jadad scale includes 5 questions to evaluate the quality of RCTs. Each question is given a maximum score of 1 point. The questions include qualifiers that must be met to obtain a point: (1) Was the trial described as randomized? (The words random, randomly, or randomization must be used.) (2) Is the method of randomization appropriate? (The methods used to generate the sequence of randomization must be described.) (3) Was the study described as double-blind? (The words double-blind must be used.) (4) Is the method of blinding appropriate? (If not, 1 point is deducted.) (5) Is there a description of withdrawals and dropouts? (The reasons need to be included, and if there are no withdrawals, it must be stated as such.)

The scale awards 1–5 points to RCTs. RCTs with ≤ 2 points or less are considered low-quality studies, and RCTs with ≥ 3 points are considered high-quality studies.²⁶

Sensitivity Analysis

The following strategies were used for the sensitivity analysis: excluding unpublished RCTs, excluding RCTs

Table 2. Inclusion and Exclusion Criteria in Studies Included in This Review

Study	Inclusion Criteria	Exclusion Criteria
Almarakbi et al ²⁷	ASA II, 18–60 y old, BMI > 30 kg/m ² , elective laparoscopic banding surgery	Asthma, COPD, restrictive lung disease, increased intracranial pressure, smoking history
Pang et al ¹⁴	ASA I–II, 16–70 y old, elective laparoscopic cholecystectomy	Pre-existing cardiac and respiratory disease requiring treatment, spontaneous pneumothorax history, evidence of hemodynamic insufficiency
Severgnini et al ²⁸	Elective non-laparoscopic abdominal surgery under general anesthesia expected to last > 2 h, > 18 y old	BMI > 40 kg/m ² ; laparoscopic or emergency surgery; previous lung surgery; persistent hemodynamic instability; intractable shock considered unsuitable for the study by the subject's managing physician; COPD history; repeated systemic corticosteroid therapy for exacerbations of COPD, asthma, or sleep disorders; recent immunosuppressive medication defined as need for chemotherapy or radiation therapy; < 2 months after chemotherapy or radiation therapy; severe cardiac disease defined as New York Heart Association class III or IV or acute coronary syndrome; persistent ventricular tachyarrhythmias; pregnancy (by lab testing); acute lung injury or ARDS, expected to require prolonged postoperative mechanical ventilation; any neuromuscular disease; contraindications to position an epidural catheter because of major clotting disorders or sign of infection at the site of the procedure
Sprung et al ²⁹	BMI > 40 kg/m ² , open bariatric surgery	Abnormalities in spirometry (FEV ₁ < 50% of predicted, FVC < 50% of predicted), active asthma (requiring bronchodilator therapy), previous lung surgery, home oxygen therapy
Weingarten et al ²⁰	> 65 y old, major open abdominal surgery	Significant pulmonary disease with abnormalities in spirometry consistent with either obstructive or restrictive pulmonary disease, active asthma (requiring chronic bronchodilator therapy), previous lung surgery, home oxygen therapy, significant cardiac dysfunction (left ventricular ejection fraction < 40%), BMI > 35 kg/m ²
Whalen et al ¹⁹	ASA II/II/III, 25–65 y old, BMI > 40 kg/m ² , laparoscopic bariatric Roux-en-Y operations	No significant preoperative pulmonary disease (FEV ₁ < 50% of predicted, FVC < 50% of predicted), active asthma, home oxygen therapy

ASA = American Society of Anesthesiologists
BMI = body mass index

published as abstracts, and excluding RCTs of lower quality as assessed by the Jadad scale (score of < 3).

Results

Results of Search

See Table 2 for inclusion and exclusion criteria and Table 3 for demographic variables. Of the 439 studies originally identified, 12 studies that met the inclusion criteria were retrieved for further scrutiny. Agreement between reviewers was achieved in all but 2 studies, which were subsequently sent to the third reviewer for further evaluation. Only one of these 2 studies was included, for a total of 6 studies receiving a Jadad score of ≥ 3 .^{14,19,20,27-29} The 2013 study by Futier et al³⁰ was not included because of the excessively large V_T in the control group (10–12 mL/kg), as well as the lack of a protocol to account for confounding variables with their primary outcome measurements.

Included Studies

Interventions. The primary intervention in all 6 studies was the utilization of alveolar recruitment maneuvers. The method for performing an alveolar recruitment maneuver varied among studies. In addition, the amount of PEEP and V_T differed between control and intervention groups among the studies (Table 4). For example, 3 studies compared subjects with higher V_T and no PEEP with subjects with lower V_T , alveolar recruitment maneuvers, and PEEP.^{14,20,28} The frequency of the alveolar recruitment maneuvers also varied depending upon the study.

Subjects. Subjects in all studies underwent abdominal surgery. Three studies involved open procedures,^{20,28,29} and 3 studies involved laparoscopic surgery.^{14,19,27} The subjects in all 3 laparoscopic studies were rated as American Society of Anesthesiologists classification I–III.^{14,19,27} Subjects in 5 studies had an average body mass index of > 25 kg/m².^{19,20,27-29} Two studies included subjects with a

Table 3. Demographic Variables in Studies Included in This Review

Study	Age (y)		Males/Females (n)		ASA Score		Anesthesia Time (min)		BMI (kg/m ²)	
	Control	Recruitment Maneuver	Control	Recruitment Maneuver	Control	Recruitment Maneuver	Control	Recruitment Maneuver	Control	Recruitment Maneuver
Almarakbi et al ²⁷	38 ± 3	38 ± 3 38 ± 3 38 ± 4	8/7	9/6 7/8 8/7	Not given	Not given	93	91 95 93	33 ± 2	33 ± 1 34 ± 1 34 ± 1
Pang et al ¹⁴	52.16 ± 13.63	49.14 ± 13.91	3/9	6/6	I/II (10/2)	I/II (9/3)	60 ± 16 (operation)	70.7 ± 28 (time)	Not given	Not given
Severgnini et al ²⁸	67 ± 9	65.5 ± 11.4	16/11	18/10	I/II/III (4/21/2)	I/II/III (6/19/3)	223 ± 80	193 ± 64	Average weight 55 ± 21 kg (21)	Average weight 56 ± 8 kg 25 ± 4.9
Sprung et al ²⁹	48 ± 9	52 ± 9	4/5	4/4	Not given	Not given	433 ± 137	458 ± 177	51 ± 5	56 ± 11
Weingarten et al ²⁰	72.1	73.8	16/4	15/5	I/II (6) III/IV (14)	I/II (9) III/IV (11)	344 ± 103	308 ± 112	27.9 ± 4.4	27.8 ± 4.3
Whalen et al ¹⁹	38 ± 11	44 ± 9	4/6	2/8	Not given	Not given	231 ± 91 (operation)	185 ± 33 (time)	53 ± 11	48 ± 6

Values are expressed as mean ± SD unless indicated otherwise.

ASA = American Society of Anesthesiologists

BMI = body mass index

body mass index > 50 kg/m².^{19,29} In 5 studies, the average age of subjects was > 40 y. Two studies included subjects with a mean age of > 65 y.^{14,20,28,29} The ratio between men and women varied with each study, but overall, there was a slightly higher number of male subjects. Specific exclusion criteria are shown in Table 2. All studies excluded subjects with significant pre-existing lung disease.

Duration of Studies. Most studies finished the majority of their outcome measures in the immediate postoperative period. Although postoperative protocols were not delineated, 4 of the 6 studies included hospital stay as a secondary outcome measure.^{19,20,27,28} The time to hospital discharge varied from days to weeks.

Risk of Bias of Included Studies

All studies were RCTs that received a Jadad score of ≥ 3. All of the study reports included a description of the randomization technique. Three of the studies included an assessment of blinding.^{19,27,28} Whalen et al¹⁹ also included blinding by the researcher collecting postoperative data. There were no dropouts in 4 studies.^{14,19,20,27} In the other 2 studies, the researchers explained the reason for dropouts. Neither of these studies included an intention-to-treat analysis.^{28,29}

Primary Outcomes

See Table 5 for primary and secondary outcomes of included studies.

Intraoperative P_{aO₂}. There was an increase in intraoperative P_{aO₂} in the alveolar recruitment maneuver groups compared with control groups in the 4 studies in which this outcome was measured.^{14,19,20,27} Sprung et al²⁹ included P_{aO₂}/F_{IO₂} in their results, whereas Severgnini et al²⁸ included only intraoperative S_{pO₂}.

Pulmonary Compliance. There was an increase in pulmonary compliance in the alveolar recruitment maneuver groups compared with the control groups in 4 of the 5 studies in which this outcome was measured.^{19,20,27,29} Severgnini et al²⁸ found no difference in pulmonary compliance among groups, whereas Pang et al¹⁴ did not include this outcome measure in their study.

P_{(A-a)O₂}. None of the studies included P_{(A-a)O₂} measurements in their results.

P_{aO₂}/F_{IO₂}. There was an increase in P_{aO₂}/F_{IO₂} in the alveolar recruitment maneuver groups compared with the control groups in the 3 studies that included this outcome measure.^{19,20,29}

Secondary Outcomes

Postoperative Pulmonary Complications. Four studies found no difference in the occurrence of postoperative pulmonary complications. Almarakbi et al²⁷ found that subjects in the alveolar recruitment maneuver group with PEEP had the shortest hospital stays. Severgnini et al²⁸ reported a higher occurrence of postoperative pulmonary

Table 4. Description of Intervention and Control Groups

Study	Control Group	Intervention Group
Almarakbi et al ²⁷	PEEP = 10 cm/H ₂ O (<i>n</i> = 15)	Experimental groups (<i>n</i> = 15 in each group): (1) Single alveolar recruitment maneuver group without PEEP: PIP = 40 cm/H ₂ O held for 15 s (2) Single alveolar recruitment maneuver group with PEEP: PIP = 40 cm/H ₂ O held for 15 s, then PEEP = 10 cm/H ₂ O for rest of surgery (3) Repeated alveolar recruitment maneuver group with PEEP: PIP = 40 cm/H ₂ O held for 15 s, then PEEP = 10 cm/H ₂ O, repeated every 10 min
Pang et al ¹⁴	V _T = 10 mL/kg, PEEP = 0 (<i>n</i> = 12)	Following intubation, manual inflation to PIP = 40 cm/H ₂ O without PEEP for 10 breaths for 1 min, then placed back on ventilator with PEEP = 5 cm/H ₂ O and V _T = 10 mL/kg (<i>n</i> = 12)
Severgnini et al ²⁸	V _T = 9 mL/kg of IBW, PEEP = 0 (<i>n</i> = 27)	V _T = 7 mL/kg of IBW, PEEP = 10 cm H ₂ O, alveolar recruitment maneuver performed after intubation, after any disconnect, and before extubation, provided stable vital signs; alveolar recruitment maneuver defined as setting PIP at 45 cm H ₂ O, I:E = 3:1, then V _T increased in steps of 4 mL/kg of IBW until plateau pressure = 30 cm H ₂ O, then 3 breaths, then returned to baseline (<i>n</i> = 28)
Sprung et al ²⁹	V _T = 8 mL/kg, PEEP = 4 (<i>n</i> = 9)	Stepwise increase in PEEP from 4 to 10 cm H ₂ O for 3 breaths, 10 to 15 cm H ₂ O for 3 breaths, 15 to 20 cm H ₂ O for 10 breaths, with maximum PIP = 50 cm H ₂ O; PEEP then set at 12 cm H ₂ O; alveolar recruitment maneuvers repeated 30 and 60 min after first recruitment (following intubation and placement of arterial line), then hourly (<i>n</i> = 8)
Weingarten et al ²⁰	V _T = 10 mL/kg of IBW, PEEP = 0 (intrinsic PEEP = 2.5 cm H ₂ O) (<i>n</i> = 20)	V _T = 6 mL/kg of IBW, PEEP = 4 cm H ₂ O, alveolar recruitment maneuver performed after intubation (exact time not given), 30 and 60 min after the first alveolar recruitment maneuver, hourly thereafter; alveolar recruitment maneuver defined as incrementally increasing PEEP from 4 to 10 cm H ₂ O for 3 breaths, 15 cm H ₂ O for 3 breaths, then 20 cm H ₂ O PEEP for 10 breaths; after alveolar recruitment maneuver, PEEP set to 12 cm H ₂ O, not stated if subsequent recruitment maneuvers decreased to 4 cm H ₂ O to start (<i>n</i> = 20)
Whalen et al ¹⁹	V _T = 8 mL/kg of IBW, PEEP = 4 (<i>n</i> = 10)	V _T = 8 mL/kg of IBW, PEEP = 4 cm H ₂ O, after pneumoperitoneum, PEEP increased in a stepwise manner from 4 to 10 cm H ₂ O for 3 breaths, 15 cm H ₂ O for 3 breaths, 20 cm H ₂ O for 10 breaths, with PEEP = 12 cm H ₂ O; alveolar recruitment maneuvers repeated until a baseline P _{aO₂} was recorded if P _{aO₂} kept increasing after the first series of alveolar recruitment maneuvers; an alveolar recruitment maneuver also instituted anytime the P _{aO₂} decreased by > 25 mm Hg of the baseline value (<i>n</i> = 10)

V_T = tidal volume

IBW = ideal body weight

PIP = peak inspiratory pressure

I:E = inspiratory-expiratory ratio

complications in the control group on postoperative day 1 compared with the alveolar recruitment maneuver group, but they found no significant difference thereafter.

Postoperative P_{aO₂} or S_{pO₂}. Four studies found no significant difference in postoperative P_{aO₂} or S_{pO₂} between groups. Almarakbi et al²⁷ found that the highest S_{pO₂} occurred in the alveolar recruitment maneuver group with PEEP. Severgnini et al²⁸ reported a statistically significant decrease in S_{pO₂} in the control group on postoperative days 1 and 3, whereas the S_{pO₂} remained the same in the alveolar recruitment maneuver group.

Complications of Alveolar Recruitment Maneuvers. Four of the studies reported no complications with the use of alveolar recruitment maneuvers. Complications of al-

veolar recruitment maneuvers were variably defined as hypotension (mean arterial blood pressure of < 60 mm Hg), hypertension (systolic blood pressure of > 150 mm Hg), an S_{pO₂} of < 90%, pneumothorax requiring a chest tube, a heart rate of < 60 beats/min, and a need to give a fluid bolus or vasoactive medication. Severgnini et al²⁸ noted comparable complications between the alveolar recruitment maneuver and control groups. Whalen et al¹⁹ identified an increase in the amount of vasopressors used in the alveolar recruitment maneuver group compared with the control group, but this difference was not statistically significant.

Frequency of Alveolar Recruitment Maneuvers. As stated earlier, the frequency of alveolar recruitment maneuvers varied among the studies in this review. Pang

ALVEOLAR RECRUITMENT MANEUVERS UNDER GENERAL ANESTHESIA

Table 5. Comparing Primary and Secondary Outcome Measures

Study	Primary Outcomes of Systematic Review				Secondary Outcomes of Systematic Review				
	Intraoperative P_{aO_2}	Pulmonary Compliance	$P_{(A-a)O_2}$	P_{aO_2}/F_{IO_2}	Postoperative Pulmonary Complications	Postoperative P_{aO_2} or S_{pO_2}	Complications	Frequency	Airway Resistance
Almarakbi et al ²⁷	Increased in single and repeated alveolar recruitment maneuvers groups with PEEP only	Increased in single and repeated alveolar recruitment maneuvers groups with PEEP only	Not studied	Not studied	Subjects in the repeated alveolar recruitment maneuver group with PEEP had the shortest hospital stay, followed by the single alveolar recruitment maneuver group with PEEP.	Highest S_{pO_2} in the repeated alveolar recruitment maneuver group with PEEP, then single alveolar recruitment maneuver group with PEEP	None	Single alveolar recruitment maneuver group with PEEP once, repeated alveolar recruitment maneuver group with PEEP every 10 min	Not studied
Pang et al ¹⁴	Increased in the alveolar recruitment maneuver group	Not studied	Not studied	Not studied	Not studied	No difference	None	Once	Not studied
Severgnini et al ²⁸	Not studied	No difference	Not studied	Not studied	Higher in the control group on postoperative day 1, no difference in hospital stay	Decrease in S_{pO_2} on days 1 and 3 in control group, remained the same in the alveolar recruitment maneuver group	Same as in the control group	Varied, no specifics given	Not studied
Sprung et al ²⁹	Not studied	Higher in the alveolar recruitment maneuver group	Not studied	Higher in the alveolar recruitment maneuver group	Not studied	No significant difference	None	After intubation, 30 and 60 min after intubation, then every hour after	Decreased in the alveolar recruitment maneuver group
Weingarten et al ²⁰	Increased in the alveolar recruitment maneuver group only	Increased in the alveolar recruitment maneuver group only	Not studied	Higher in the alveolar recruitment maneuver group	No difference in hospital stay	No difference	None	After intubation, 30 and 60 min after intubation, then every hour after	Decreased in the alveolar recruitment maneuver group
Whalen et al ¹⁹	Increased more in the alveolar recruitment maneuver group	Higher in the alveolar recruitment maneuver group initially and after insufflation	Not studied	Higher in the alveolar recruitment maneuver group	No difference	No difference	None	Varied, see study	No difference

 $P_{(A-a)O_2}$ = alveolar-arterial oxygen difference

et al¹⁴ utilized an alveolar recruitment maneuver only once, whereas Sprung et al²⁹ and Weingarten et al²⁰ performed alveolar recruitment maneuvers after intubation, 30 and 60 min following intubation, and every hour thereafter. The study by Almarakbi et al²⁷ was unique in that it was the only study to include the effects of different frequencies of alveolar recruitment maneuvers. These researchers compared alveolar recruitment maneuvers performed once with those performed every 10 min.

Discussion

Summary of Main Results

This review of alveolar recruitment maneuvers included 6 RCTs, all of which achieved a score of ≥ 3 on the Jadad scale. The studies differed in their subjects, independent/dependent variables, control and intervention groups, technique of applying an alveolar recruitment maneuver, and outcome measures, making absolute quantitative comparisons treacherous. We included the major clinical themes section below to qualitatively approach comparisons in this systematic review. Despite differences in methods, certain comparisons can be made among the 6 studies. Subjects in all studies underwent abdominal surgery. Most subjects had a body mass index of $> 25 \text{ kg/m}^2$ and no significant pre-existing lung disease. Most studies utilized one of 2 methods for the application of alveolar recruitment maneuvers, with the exception of Severgnini et al.²⁸ Alveolar recruitment maneuvers consisted of either a step-wise increase in PEEP with sustained breaths or sustained manual inflations of an anesthesia reservoir bag to a PIP of 40 cm H₂O. All studies reported $P < .05$ as a statistically significant outcome difference.

Primary Outcome Measures

Overall, subjects in the alveolar recruitment maneuver groups experienced a higher intraoperative P_{aO_2} , greater lung compliance, and higher P_{aO_2}/F_{IO_2} .

Secondary Outcome Measures

Overall, there were no significant differences in the rate of postoperative pulmonary complications, postoperative P_{aO_2} , S_{pO_2} , or intraoperative airway resistance between subjects in the alveolar recruitment maneuver and control groups in the 6 studies in this review. Despite the variety of maximum PIP allowed during alveolar recruitment maneuvers in the studies (ranging from 40 to 50 cm H₂O), the occurrence of complications associated with alveolar recruitment maneuvers appeared minimal to nonexistent. Five of the 6 studies reported no complications associated with the use of alveolar recruitment maneuvers. The study that

reported complications in the alveolar recruitment maneuver group had similar complications in the control group.²⁸ The frequency of alveolar recruitment maneuvers varied among studies in this review.^{14,19,20,27-29}

Major Clinical Themes

Is There Evidence to Support Alveolar Recruitment Maneuvers in Open Abdominal Surgery? Of the 3 studies involving open abdominal surgery, only Weingarten et al²⁰ included the effects of alveolar recruitment maneuvers on intraoperative P_{aO_2} , which increased solely in the alveolar recruitment maneuver group. Sprung et al²⁹ and Weingarten et al²⁰ reported an increase in pulmonary compliance along with an increase in P_{aO_2}/F_{IO_2} in the alveolar recruitment maneuver groups. Severgnini et al²⁸ found no significant difference in pulmonary compliance among groups and did not report on P_{aO_2}/F_{IO_2} . None of the studies reported statistically significant complications during or after alveolar recruitment maneuvers. There was no consensus on the benefit of alveolar recruitment maneuvers in the postoperative period.

Is There Evidence to Support Alveolar Recruitment Maneuvers in Laparoscopic Abdominal Surgery? All 3 studies involving laparoscopic abdominal surgery showed an increase in intraoperative P_{aO_2} with alveolar recruitment maneuvers.^{14,19,27} Both Almarakbi et al²⁷ and Whalen et al¹⁹ also reported an increase in pulmonary compliance, whereas Pang et al¹⁴ did not include pulmonary compliance in their research. Only Whalen et al¹⁹ reported on intraoperative P_{aO_2}/F_{IO_2} , which was increased in the alveolar recruitment maneuver group. There was no consensus of evidence regarding postoperative benefits of alveolar recruitment maneuvers. Almarakbi et al²⁷ reported a higher S_{pO_2} and shorter hospital stays with the alveolar recruitment maneuver groups, whereas Whalen et al¹⁹ found no differences in the postoperative period between groups. The lack of consensus for the benefits of alveolar recruitment maneuvers extending into the postoperative period will be discussed below in the implications for practice section.

Which Alveolar Recruitment Maneuver Is the Best?

Excluding the study by Severgnini et al,²⁸ the different types of alveolar recruitment maneuvers were found to be equally effective. Both Almarakbi et al²⁷ and Pang et al¹⁴ utilized sustained manual inflations up to a PIP of 40 cm H₂O for their alveolar recruitment maneuvers. Almarakbi et al²⁷ utilized single sustained manual inflations for 15 s, whereas Pang et al¹⁴ utilized 10 sustained manual inflations over 1 min. Both studies found an increase in intraoperative P_{aO_2} in the alveolar recruitment maneuver groups. However, only Almarakbi et al²⁷ reported on pulmonary compliance, which showed an increase exclusively

in the alveolar recruitment maneuver groups. Neither study reported on P_{aO_2}/F_{IO_2} or airway resistance.

Three studies utilized stepwise increases in PEEP starting at 4 cm H₂O and ending at 20 cm H₂O as alveolar recruitment maneuvers.^{19,20,29} Of the 2 studies that included intraoperative P_{aO_2} ,^{19,20} there was a statistically significant increase in the alveolar recruitment maneuver groups. All 3 studies showed an improvement in pulmonary compliance in the alveolar recruitment maneuver groups. Only Sprung et al²⁹ and Whalen et al¹⁹ included P_{aO_2}/F_{IO_2} as an outcome measure, which was significantly increased in the alveolar recruitment maneuver groups in the intraoperative period.

Manual sustained hyperinflations require a shorter amount of time to perform, whereas stepwise increases in PEEP or V_T may prevent untoward subject responses such as straining or coughing when the depth of anesthesia is equivocal.

How Long Should Alveolar Recruitment Maneuvers Be Held to Be Effective?

The time that the manual alveolar recruitment maneuver is held to a set PIP varies in the literature. As stated previously, some authors consider inflation of the lungs for 5–30 s to a prescribed PIP as representative of alveolar recruitment maneuvers.¹⁵⁻¹⁸ Almarakbi et al²⁷ held alveolar recruitment maneuvers for 15 s. Pang et al¹⁴ did not specify a period of time for holding the alveolar recruitment maneuver. They utilized 10 alveolar recruitment maneuvers over 1 min, which leaves, at best, a few seconds for an inspiratory hold if it indeed was utilized. The 3 studies involving stepwise increases in PEEP all performed them in the same manner.^{19,20,29} Subjects received 3 breaths when the PEEP was increased from 4 to 10 cm H₂O, then received 3 breaths when the PEEP was increased from 10 to 15 cm H₂O, and finally 10 breaths when the PEEP was increased to 20 cm H₂O. Severgnini et al²⁸ defined an alveolar recruitment maneuver as setting PIP to 45 cm H₂O with an inspiratory/expiratory ratio of 3:1. The V_T was then increased in steps of 4 mL/kg of ideal body weight until a plateau pressure of 30 cm H₂O was reached, at which time 3 breaths were given.

Is There Evidence to Support the Use of Repeated Alveolar Recruitment Maneuvers?

The repeated-measures alveolar recruitment maneuver group in the study by Almarakbi et al²⁷ showed the greatest improvement in intraoperative P_{aO_2} , pulmonary compliance, and postoperative P_{aO_2} . Most of the studies repeated alveolar recruitment maneuvers at various times throughout the surgery. The study by Almarakbi et al²⁷ was the only study to measure the effects of different frequencies of alveolar recruitment maneuvers. One of the experimental groups received an alveolar recruitment maneuver once, whereas

another received alveolar recruitment maneuvers every 10 min until completion of surgery. The researchers noted that the benefit of an elevated P_{aO_2} in the single alveolar recruitment maneuver group was temporary, whereas the benefit was sustained in the repeated-measures alveolar recruitment maneuver group. The subjects in the repeated-measures alveolar recruitment maneuver group also had the shortest hospital stay.

Is There Evidence to Support Using Alveolar Recruitment Maneuvers Followed By PEEP?

The only study to isolate the differences in the utilization of PEEP following an alveolar recruitment maneuver was undertaken by Almarakbi et al.²⁷ As stated above, these researchers compared the effects of repeated alveolar recruitment maneuvers versus a single alveolar recruitment maneuver. They also included a third experimental group that received a single alveolar recruitment maneuver followed by zero PEEP. There were no significant differences between the single alveolar recruitment maneuver group without PEEP and the control group (PEEP of 10 cm H₂O only). Neither group showed an improvement in respiratory compliance, intraoperative or postoperative P_{aO_2} , or a reduction in hospital stay. There was, however, a significant advantage in compliance, intraoperative and postoperative P_{aO_2} , and reduction in hospital stay in the alveolar recruitment maneuver groups followed by PEEP.

What Is the Best Method for Determining the Effectiveness of Alveolar Recruitment Maneuvers?

Besides obtaining an arterial blood gas and calculating the $P_{(A-a)O_2}$, there is a more practical and less invasive approach. Four of the 5 studies found a statistically significant increase in pulmonary compliance in alveolar recruitment maneuver groups.^{19,20,27,29} Dynamic pulmonary compliance is calculated by dividing V_T by the difference between PIP and PEEP: dynamic compliance = $V_T/(PIP - PEEP)$. Static pulmonary compliance is calculated by dividing V_T by the difference between the plateau pressure (P_{plat}) and PEEP: static compliance = $V_T/(P_{plat} - PEEP)$. An increase in pulmonary compliance immediately following the application of an alveolar recruitment maneuver would presumably reflect a reduction in atelectasis and result in an increase in P_{aO_2} due to better matching of ventilation and perfusion. To avoid the need to perform the above calculations, a more clinically relevant method for determining whether an increase in compliance has occurred can be utilized. During volume control continuous mandatory ventilation, the provider would note that the same V_T is delivered at a lower PIP following application of alveolar recruitment maneuvers. With pressure control continuous mandatory ventilation, the provider would note that an increased V_T is delivered at the same inspiratory pressure setting. Contemporary anesthesia ventilators often include options to

display the pressure-volume loop and calculated dynamic compliance values, making it even easier to measure the effectiveness of alveolar recruitment maneuvers.

Overall Completeness and Applicability of Evidence

Evidence from the 6 included studies regarding the benefit of alveolar recruitment maneuvers in the postoperative period was lacking.^{14,19,20,27-29} Although postoperative measurements were evaluated, no study included a postoperative pulmonary protocol accounting for confounding variables. In addition, only one study included dosages used for reversal of neuromuscular blockade.¹⁴ Other primary and secondary outcomes were compared as noted. The applicability of the evidence summarized in this review is limited to adult subjects without extensive pre-existing pulmonary disease who are undergoing abdominal surgery.

Quality of Evidence

This systematic review was limited to RCTs, systematic reviews, and quantitative meta-analyses. Only high-quality RCTs that matched the inclusion criteria were selected. Of concern, the sample sizes of the included studies were small, which brings into question the ability of the studies to control and account for confounding variables. This statistical fragility may have elevated the risk of type I or type II errors. The lack of specific emergence or postoperative pulmonary protocols limits the applicability of postoperative measurements.

Potential Bias in the Review Process

The reviewers did not examine the results of individual studies until after all studies met the inclusion criteria, including achieving a score of ≥ 3 on the Jadad scale. Reviewers were therefore unaware of the content of the included studies as it related to the benefit of alveolar recruitment maneuvers until after the studies were selected.

Conclusions

Implications for Practice

Depending on the type of surgery and patient status, the practitioner should utilize alveolar recruitment maneuvers followed by PEEP after induction of general endotracheal anesthesia, routinely, and in the presence of a decreasing S_{pO_2} . Although evidence from this review limits the application of alveolar recruitment maneuvers to adults undergoing abdominal surgery, there is a wealth of literature to support alveolar recruitment maneuvers in other surgeries.³¹⁻³⁴ Teleologically, a healthy human sighs an average of 9–10 times/h. Thus, most patients undergoing general endotracheal anes-

thesia would benefit from this homeostatic reflex in the form of an alveolar recruitment maneuver.

Compared with the control group, all types of alveolar recruitment maneuvers were beneficial in the intraoperative period. Manual sustained hyperinflations require a short amount of time to perform, whereas stepwise increases in PEEP or V_T may prevent untoward subject responses such as coughing or straining when the depth of anesthesia is questionable. Alveolar recruitment maneuvers should be performed instead of solely increasing F_{IO_2} during the maintenance phase of anesthesia. Continually increasing F_{IO_2} in the face of a decreasing S_{pO_2} may maintain an elevated S_{pO_2} , but may also mask a physiological shunt. Oxygen can also adversely affect respiratory control, ventilation/perfusion ratios, and hypoxic pulmonary vasoconstriction, as well as cause vasoconstriction of systemic arterioles.³⁵ Oxygen products from reduced nicotinamide adenine dinucleotide phosphate oxidase also increase inflammation and tissue injury through a complex interaction among neutrophils, macrophages, and platelets.³⁶ Alveolar recruitment maneuvers allow the anesthesia provider to reduce the F_{IO_2} yet maintain a higher S_{pO_2} .

Current evidence does not support the assumption that the benefit of alveolar recruitment maneuvers extends to the postoperative period because none of the studies included a specific emergence or postoperative pulmonary protocol. Emergence from anesthesia and the immediate postoperative period can leave a patient susceptible to atelectasis and hypoxemia. Patients often breathe low V_T without PEEP and on 100% oxygen for minutes during this time. Given the negative consequences of absorption atelectasis, low V_T , and the absence of PEEP, techniques for emergence need to be re-examined to prevent the negation of intraoperative alveolar recruitment maneuvers. Teleologically, it seems implicit that reducing intraoperative risk may decrease postoperative issues.

Implications for Research

Most studies involve the use of alveolar recruitment maneuvers during the maintenance phase of anesthesia. To maximize the benefits of alveolar recruitment maneuvers for patients in the postoperative period, a study incorporating sigh breaths following extubation with the utilization of CPAP should be undertaken. Sigh breaths in the postoperative period can be quantified with the use of an incentive spirometer. After extubation, patients control their own gas composition and may benefit from CPAP in the post-anesthesia care unit to prevent atelectasis secondary to diminished V_T . The use of CPAP in the postoperative period significantly decreases the incidence of pneumonia, sepsis, and hypoxemia.³⁷ A study on the use of intraoperative alveolar recruitment maneuvers combined with the use of CPAP and sigh breaths following extubation may

extend the benefits of alveolar recruitment maneuvers into the recovery period, reducing postoperative pulmonary complications and improving subject outcomes.

Patient position during emergence and following extubation may also improve patient outcomes and reduce atelectasis formation. A study comparing elevated head-of-bed and supine positions during emergence and in the postoperative period may show a reduction in atelectasis and improvement in oxygenation.

REFERENCES

- Bendixen HH, Smith GM, Mead J. Pattern of ventilation in young adults. *J Appl Physiol* 1964;19:195-198.
- Bartlett D Jr. Origin and regulation of spontaneous deep breaths. *Respir Physiol* 1971;12(2):230-238.
- Ferris BG Jr, Pollard DS. The effect of deep and quiet breathing on pulmonary compliance in man. *J Clin Invest* 1960;39:143-149.
- Bendixen HH, Bullwinkel B, Hedley-Whyte J, Laver MB. Atelectasis and shunting during spontaneous ventilation in anesthetized subjects. *Anesthesiology* 1964;25:297-301.
- Hedenstierna G, Edmark L. Mechanisms of atelectasis in the perioperative period. *Best Pract Res Clin Anaesthesiol* 2010;24(2):157-169.
- Dietl P, Frick M, Mair N, Bertocchi C, Haller T. Pulmonary consequences of a deep breath revisited. *Biol Neonate* 2004;85(4):299-304.
- Froese AB, Bryan AC. Effects of anesthesia and paralysis on diaphragmatic mechanics in man. *Anesthesiology* 1974;41(3):245-255.
- Tokics L, Hedenstierna G, Strandberg A, Brismar B, Lundquist H. Lung collapse and gas exchange during general anesthesia: effects of spontaneous breathing, muscle paralysis, and positive end-expiratory pressure. *Anesthesiology* 1987;66(2):157-167.
- Magnusson L, Spahn DR. New concepts of atelectasis during general anaesthesia. *Br J Anaesth* 2003;91(1):61-72.
- Brismar B, Hedenstierna G, Lundquist H, Strandberg A, Svensson L, Tokics L. Pulmonary densities during anesthesia with muscular relaxation—a proposal of atelectasis. *Anesthesiology* 1985;62(4):422-428.
- Joyce CJ, Baker AB. What is the role of absorption atelectasis in the genesis of perioperative pulmonary collapse. *Anaesth Intensive Care* 1995;23(6):691-696.
- Hedenstierna G, Rothen HU. Atelectasis formation during anesthesia: causes and measures to prevent it. *J Clin Monit Comput* 2000;16(5-6):329-335.
- Gonçalves LO, Cicarelli DD. Alveolar recruitment maneuver in anesthetic practice: how, when, and why it may be useful. *Rev Bras Anesthesiol* 2005;55(6):631-638.
- Pang CK, Yap J, Chen PP. The effect of an alveolar recruitment strategy on oxygenation during laparoscopic cholecystectomy. *Anaesth Intensive Care* 2003;31(2):176-180.
- Biddle C, Epps L, Hassanein R. The efficacy of sighs in subjects with COPD undergoing general anesthesia and controlled ventilation. *AANA J* 1989;57(2):131-136.
- Cakmakaya OS, Kaya G, Altintas F, Hayirlioglu M, Ekici B. Restoration of pulmonary compliance after laparoscopic surgery using a simple alveolar recruitment maneuver. *J Clin Anesth* 2009;21(6):422-426.
- Celebi S, Köner O, Menda F, Korkut K, Suzer K, Cakar N. The pulmonary and hemodynamic effects of two different recruitment maneuvers after cardiac surgery. *Anesth Analg* 2007;104(2):384-390.
- Rothen HU, Neumann P, Berglund JE, Valtysson J, Magnusson A, Hedenstierna G. Dynamics of re-expansion of atelectasis during general anaesthesia. *Br J Anaesth* 1999;82(4):551-556.
- Whalen FX, Gajic O, Thompson GB, Kendrick ML, Que FL, Williams BA, et al. The effects of the alveolar recruitment maneuver and positive end-expiratory pressure on arterial oxygenation during laparoscopic bariatric surgery. *Anesth Analg* 2006;102(1):298-305.
- Weingarten TN, Whalen FX, Warner DO, Gajic O, Schears GJ, Snyder MR, et al. Comparison of two ventilatory strategies in elderly subjects undergoing major abdominal surgery. *Br J Anaesth* 2010;104(1):16-22.
- Shander A, Fleisher LA, Barie PS, Bigatello LM, Sladen RN, Watson CB. Clinical and economic burden of postoperative pulmonary complications: subject safety summit on definition, risk-reducing interventions, and preventive strategies. *Crit Care Med* 2011;39(9):2163-2172.
- Tusman G, Böhm SH, Warner DO, Sprung J. Atelectasis and perioperative pulmonary complications in high-risk subjects. *Curr Opin Anaesthesiol* 2012;25(1):1-10.
- Magnusson L, Zemgulis V, Wicky S, Tydén H, Thelin S, Hedenstierna G. Atelectasis is a major cause of hypoxemia and shunt after cardiopulmonary bypass: an experimental study. *Anesthesiology* 1997;87(5):1153-1163.
- Linde-Zwirble WL, Bloom JD, Mecca RS, Hansell DM. Postoperative pulmonary complications in adult elective surgery subjects in the US: severity, outcomes, and resources use. *Critical Care* 2010;14(Suppl 1):P210.
- Paulus F, Binnekade JM, Vroom MB, Schultz MJ. Benefits and risks of manual hyperinflation in intubated and mechanically ventilated intensive care unit subjects: a systematic review. *Crit Care* 2012;16(4):R145.
- Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gaghaghian DJ, McQuay HJ. Assessing the quality of reports of randomized clinical trials: Is blinding necessary? *Controlled Clin Trials* 1996;17(1):1-12.
- Almarakbi WA, Fawzi HM, Alhashemi JA. Effects of four intraoperative ventilatory strategies on respiratory compliance and gas exchange during laparoscopic gastric banding in obese subjects. *Br J Anaesth* 2009;102(6):862-868.
- Severgnini P, Selmo G, Lanza C, Chiesa A, Frigerio A, Bacuzzi A, et al. Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *Anesthesiology* 2013;118(6):1307-1321.
- Sprung J, Whalen FX, Comfere T, Bosnjak ZJ, Bajzer Z, Gajic O, et al. Alveolar recruitment and arterial desflurane concentration during bariatric surgery. *Anesth Analg* 2009;108(1):120-127.
- Futier E, Constantin JM, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013;369(5):428-437.
- Duff JP, Rosychuk RJ, Joffe AR. The safety and efficacy of sustained inflations as a lung recruitment maneuver in pediatric intensive care unit patients. *Intensive Care Medicine* 2007;33(10):1778-1786.
- Fan E, Wilcox ME, Brower RG, Stewart TE, Mehta S, Lapinsky SE, et al. Recruitment maneuvers for acute lung injury: a systematic review. *Am J Respir Crit Care Med* 2008;178(11):1156-1163.
- Claxton BA, Morgan P, McKeague H, Mulpur A, Berridge J. Alveolar recruitment strategy improves arterial oxygenation after cardiopulmonary bypass. *Anaesthesia* 2003;58(2):111-116.
- Unzueta C, Tusman G, Suarez-Sipmann F, Böhm S, Moral V. Alveolar recruitment improves ventilation during thoracic surgery: a randomized controlled trial. *Br J Anaesth* 2012;108(3):517-524.
- Lumb AB, Walton LJ. Perioperative oxygen toxicity. *Anesthesiol Clin* 2012;30(4):591-605.
- Ward PA, Warren JS, Johnson KJ. Oxygen radicals, inflammation, and tissue injury. *Free Radic Biol Med* 1988;5(5-6):403-408.
- Squadron V, Cocha M, Cerutti E, Schellino MM, Biolino P, Occella P, et al. Continuous positive airway pressure for treatment of postoperative hypoxemia: a randomized controlled trial. *JAMA* 2005;293(5):589-595.