

Effects of Manual Rib Cage Compressions on Mucus Clearance in Mechanically Ventilated Pigs

Akira Ouchi, Hideaki Sakuramoto, Takeshi Unoki, Yasuyo Yoshino, Haruhiko Hosino, Yasuaki Koyama, Yuki Enomoto, Nobutake Shimojo, Taro Mizutani, and Yoshiaki Inoue

BACKGROUND: Manual rib cage compression is a chest physiotherapy technique routinely used in clinical practice. However, scientific evidence remains scarce on the effects of manual rib cage compression on airway clearance and oxygenation in mechanically ventilated patients. **METHODS:** Anesthetized pigs were intubated via the trachea and mechanically ventilated. To create atelectasis, artificial mucus was infused into the airway. Each pig was randomly assigned to 1 of 2 groups: closed suctioning alone (control group, 7 pigs), or manual rib cage compression combined with closed suctioning (manual rib cage compression group, 8 pigs). Hard and brief rib cage compression synchronized with early expiratory phase was tested. Mucus clearance and oxygenation were assessed after the intervention. Sequential changes of hemodynamics were assessed after the intervention. **RESULTS:** During hard manual rib cage compression, the mean \pm SD peak expiratory flow increased to 44 ± 7 L/min compared with 31 ± 7 L/min without treatment ($P < .001$). Manual rib cage compression combined with endotracheal suctioning increased mucus clearance compared with closed suctioning alone (mucus amounts, 5.5 [3.4–9.4] g vs 0.7 [0.5–2.0] g; $P = .004$); however, it did not improve gas exchange and radiologic findings. There were no significant differences in hemodynamic variables between the 2 groups. **CONCLUSIONS:** Our findings indicated that hard and brief manual rib cage compression combined with closed suctioning was safe and led to improvement of mucus clearance; however, no effectiveness was confirmed with regard to oxygenation and ventilation. *Key words:* mechanical ventilation; airway clearance; rib cage compression; physiotherapy; suctioning. [Respir Care 0;0(0):1–9. © 0 Daedalus Enterprises]

Introduction

Retention of airway secretions can result in serious clinical complications in patients on mechanical ventilation.¹⁻⁴ Mucus retention narrows or occludes airways,^{5,6} causes breathing discomfort, and leads to atelectasis and gas exchange impairment.⁷ In the acute care setting, mucus retention is related to significantly higher mortality and morbidity.⁸ Manual rib cage

compression (MRCC) is a chest physiotherapy technique routinely used in clinical practice.⁹⁻¹¹ However, scientific evidence on the effects of MRCC on airway clearance and oxygenation in mechanically ventilated patients remains unclear because of the variety of techniques.

This work was performed at the University of Tsukuba.

This work was supported in part by a Grant-in-Aid for Young Scientists B from the Ministry of Education, Culture, Sports, Science and Technology of Japan (16K20746) and the Japan Society for the Promotion of Science.

The authors have disclosed no conflicts of interest.

Correspondence: Yoshiaki Inoue MD PhD, Department of Emergency and Critical Care Medicine, Faculty of Medicine, University of Tsukuba, Tsukuba, Ibaraki, 305-8575 Japan. E-mail: yinoue@md.tsukuba.ac.jp.

DOI: 10.4187/respcare.07249

Mr Ouchi, Mr Hosino, Dr Koyama, Dr Enomoto, Dr Shimojo, and Dr Inoue are affiliated with the Department of Emergency and Critical Care Medicine, Faculty of Medicine, University of Tsukuba, Tsukuba, Ibaraki, Japan. Mr Sakuramoto is affiliated with the Department of Adult Health Nursing, College of Nursing, Ibaraki Christian University, Hitachi, Ibaraki, Japan. Mr Unoki is affiliated with the Department of Adult Health Nursing, School of Nursing, Sapporo City University, Sapporo, Japan. Ms Yoshino is affiliated with Department of Nursing, Kanto Gakuin University College of Nursing, Yokohama, Kanagawa, Japan. Dr Mizutani is affiliated with the Ibaraki Western Medical Center, Ibaraki, Japan.

MANUAL RIB CAGE COMPRESSION IN A PORCINE MODEL

A previous animal study showed that a hard and brief MRCC technique applied during the early expiratory phase increases expiratory flow and mucus movement compared with a soft and gradual MRCC technique applied during the late expiratory phase.¹² Nevertheless, it is not clear whether MRCC combined with endotracheal suctioning promotes mucus clearance and improves oxygenation in patients with mucus retention. The objective of this study was to determine whether a combination of hard MRCC and endotracheal suctioning could improve mucus clearance and oxygenation in well-controlled mechanically ventilated pigs with retention of artificial mucus.

Methods

Animal Preparation

This study was conducted at the University of Tsukuba, Japan. The protocol was approved by the institution's animal research committee. Animals were managed according to the National Institute of Health guidelines for the Use and Care of Animals. Fifteen female pigs (mean \pm SD, 33 \pm 2 kg) were anesthetized by using midazolam and ketamine, and were restrained in a supine position. Intravenous lines were installed at the ear vein of each pig, and a continuous infusion of pentobarbital was delivered. Tracheostomy was performed with the pig under local anesthesia by using 1.0% lidocaine solution and a 7-mm inner-diameter endotracheal tube (ETT) placed in the trachea. The animals were then ventilated with LTV-1000 ventilators (CareFusion, San Diego, California).

To instill artificial mucus into the airways, a closed suctioning catheter system (Trachcare, Ballard Medical Products, Draper, Utah) was connected to the ETT. Mechanical ventilation was set as follows: pressure control, F_{IO_2} 1.0, PEEP 10 cm H₂O, peak inspiratory pressure 20 cm H₂O achieved tidal volume of 6–8 mL/kg, and breathing frequency was adjusted to maintain normocapnia. Internal ETT cuff pressure was maintained at 25 cm H₂O. The pigs were paralyzed with a loading (0.6 mg/kg) and maintenance (0.4–0.6 mg/kg/h) dose of vecuronium. Femoral artery catheter was surgically inserted to monitor the systemic arterial pressure and to collect blood samples. A pulmonary artery catheter (Model 744F8, Edwards Lifesciences, Irvine, California) was also placed via femoral vein to measure hemodynamic variables, including stroke volume and cardiac output.

Model of Atelectasis

To create atelectasis, artificial mucus was infused into the airways by using the closed suction method. The technical details are shown in our previous experimental reports.^{13,14} Briefly, the artificial mucus was made by mixing 1.6% polyethylene oxide powder and 0.1% methylene blue in a phosphate buffered saline solution, and, to prepare thinner artificial

QUICK LOOK

Current knowledge

Mechanically ventilated patients often have impaired airway mucus clearance. Manual rib cage compression is routinely used in clinical practice. However, the effect of manual rib cage compression on airway clearance and oxygenation in these patients remains unclear.

What this paper contributes to our knowledge

In this prospective randomized animal study, manual rib cage compression combined with closed suctioning resulted in a larger amount of mucus clearance than did closed suctioning alone but did not improve gas exchange.

mucus, iopamidol was mixed with the artificial mucus at a ratio of 3:2.^{13,14} Dynamic viscoelasticity of the solution was measured with a controlled shear rate rheometer, at a driving frequency of a 1 rad/s. The loss modulus (G'') of the artificial mucus was 1.6 Pa, and the storage modulus (G') was 0.97 Pa. The artificial mucus infusion was continued at a rate of 4 mL/min for 10 min (total of 48 g) with reference to a previous animal study.¹⁵ After the mucus infusion, the closed suction catheter was changed to the other catheter. The animals were maintained on mechanical ventilation for 30 min after administration of the mucus.

Hard-Type Rib Cage Compression

A single operator, a trained nurse (HH), performed the hard MRCC. The nurse applied short (\sim 1 s) and strong bilateral compressions.¹² Hand pressure pads were used to assess the force of each compression.¹⁴ In preliminary experiments, we confirmed that hand pressure reflects esophageal pressure. Compressions were synchronized with the start of the expiratory phase.

Endotracheal Suctioning

An experienced nurse (YY) performed the endotracheal suctioning according to the guidelines of the American Association for Respiratory Care.¹⁶ Neither hyperventilation nor manual hyperinflation was performed before or after endotracheal suctioning. Closed suction was performed by using the closed suctioning catheter (12 French) system mentioned above. A premeasured catheter was inserted, without applying suction, to the ETT tip. When in place, suction was applied at a pressure of -150 mm Hg for 15 s while simultaneously withdrawing the catheter. The weight of the suctioned mucus was

MANUAL RIB CAGE COMPRESSION IN A PORCINE MODEL




Fig. 1. Experimental protocol. A researcher applied a 10-min session of manual rib cage compression (MRCC). Radiologic images were taken at baseline (before randomization) and the end of the period. MI = mucus infusion.

calculated by subtracting the weight of the container before suction from its weight after suction.

Data Collection

Heart rate and mean arterial pressure were monitored by using a Philips IntelliVue MP50 Patient Monitor (Philips Medizin Systeme, Böblingen, Germany). Cardiac output was monitored by using Vigilance II (Edwards Lifesciences). Arterial blood gases were measured by using blood drawn from the femoral artery via an ABL 735 blood gas analyzer (Radiometer Copenhagen, Copenhagen, Denmark). Expiratory tidal volume, airway pressure, and airway flow were recorded by using a CP-100 pulmonary monitor (Bicore, Irvine, California). Radiologic images were taken by using a C-arm (DHF-105CX, Hitachi, Japan). Radiologic findings were assessed from the atelectasis score (0, clear lung fields; 1, plate-like atelectasis or slight infiltration; 2, partial atelectasis; 3, lobar atelectasis; and 4, bilateral lobar atelectasis)¹⁷ and 0–10 cm visual analog scale by 2 investigators blinded to all experimental information.

Study Design

The experimental protocol is illustrated in Figure 1. After the first radiologic images were taken, the pigs were randomly assigned to 2 groups: closed suctioning alone (control group, 7 pigs), or hard MRCC combined with closed suctioning (MRCC group, 8 pigs). After the experiments, the pigs were sacrificed with an intravenous injection of 10 mL (10 mEq/10 mL) potassium chloride solution.

Table 1. Baseline Characteristics (before mucus infusion)

Characteristic	Control Group	MRCC Group	<i>P</i>
Body weight, kg	33.0 ± 2.2	33.4 ± 2.8	.74
pH	7.47 ± 0.03	7.48 ± 0.02	.67
P _{aO₂} /F _{IO₂} , mm Hg	516 ± 21	500 ± 35	.33
P _{aCO₂} , mm Hg	38 ± 2	40 ± 2	.27
Dynamic compliance, mL/cm H ₂ O	27.7 ± 7.3	28.3 ± 7.2	.88
Lactate, mmol/L	2.4 ± 1.4	2.0 ± 0.6	.52
Heart rate, beats/min	135 ± 38	135 ± 29	.99
Mean arterial pressure, mm Hg	104 ± 13	114 ± 20	.26
Cardiac output, L/min	3.3 ± 0.4	3.7 ± 0.4	.77
Stroke volume, mL	28 ± 7	28 ± 7	.96

Data are mean ± SD.
MRCC = manual rib cage compression

Statistical Analysis

All values are reported as a mean ± SD unless otherwise specified. Parametric and nonparametric analyses were used in accordance with the results of the Shapiro-Wilk test. Paired *t* test or the Wilcoxon rank-sum test were used to compare before and after treatment measurements. Intra-intergroup differences were compared by using one-way analysis of variance with adjusted Bonferroni correction. Repeated measures analysis of variance was used to determine intragroup differences. Specific intergroup differences and time points of these differences were determined by using the Bonferroni correction for multiple comparisons. All tests were performed as 2-sided, with a significance level of 5%. All analyses were performed by using SPSS version 25 software (SPSS, Chicago, Illinois).

Results

Baseline characteristics of the animals in the study groups are shown in Table 1. There were no differences in body weight, hemodynamic variables, and gas exchange parameters before the artificial mucus infusion. During the compression, the mean ± SD peak expiratory flow (PEF) increased to 44 ± 7 L/min compared with 31 ± 7 L/min without treatment (*P* < .001). The mean ± SD expiratory tidal volume increased to 345 ± 46 mL compared with 215 ± 87 mL (*P* < .001) without treatment. As shown in Figure 2, MRCC combined with endotracheal suctioning increased mucus clearance compared with closed suctioning without MRCC (*P* = .004). The median range values of mucus amounts (g) in the control and MRCC groups were 0.7 (0.5–2.0) g and 5.5 (3.4–9.4) g, respectively.

The results of P_{aO₂}/F_{IO₂} and P_{aCO₂} are shown in Figure 3. With regard to oxygenation, there were no significance differences in P_{aO₂}/F_{IO₂} before mucus infusion or at baseline. After mucus infusion, P_{aO₂}/F_{IO₂} decreased in all the pigs.

MANUAL RIB CAGE COMPRESSION IN A PORCINE MODEL




Fig. 2. Sputum volume produced after control and manual rib cage compression (MRCC) interventions. Horizontal lines denote median value, upper and lower edges represent the 25th and 75th percentiles, and whiskers represent the 10th and 90th percentiles. * $P < .01$, compared with control.

Throughout the experiment, there were no significance differences in P_{aO_2}/F_{IO_2} between the control and MRCC groups (Fig. 3A). With regard to ventilation, P_{aCO_2} for both groups increased significantly compared with the baseline after the mucus infusion. There were no significance differences in P_{aCO_2} between the control and MRCC groups throughout the experiment (Fig. 3B).

Radiologic data are expressed as median (interquartile range). There were no significance differences between the groups in the atelectasis scores before and after the interventions (Fig. 4A). There were also no significance differences between the groups in visual analog scale scores before and after the interventions (Fig. 4B). The effects of MRCC on hemodynamics are shown in Table 2. There were no significance differences in heart rate, mean arterial pressure, and cardiac output between the groups.

Discussion

The purpose of the present study was to investigate whether the combination of MRCC and following closed suctioning could improve mucus clearance and oxygenation under well-controlled experimental settings. The present study demonstrated that MRCC increased the PEF and improved mucus clearance; however, the maneuver did not improve oxygenation or radiologic findings. An animal study by Marti et al¹² compared 2 different MRCC techniques and investigated their respective effects. The soft and gradual MRCC technique did not increase PEF; however, hard and brief MRCC significantly increased PEF and mucus movement.¹² MRCC generates an expiratory flow bias




Fig. 3. Changes in (A) P_{aO_2}/F_{IO_2} (P/F) and (B) P_{aCO_2} in the study groups. There were no significance differences between the 2 groups. Data are shown as means with 95% confidence intervals. MRCC = manual rib cage compression; MI = mucus infusion.

that is believed to move secretions toward the central airways through a 2-phase gas-liquid transport.^{18,19} A previous study showed that mucus clearance significantly improves by increasing PEF to 40–60 L/min.²⁰

The findings of the present study were that MRCC increased PEF, and, via the 2-phase gas-liquid transport mechanism, increased mucus clearance. Nevertheless, similar to previous findings,⁹ we did not find any positive effect on gas exchanges and radiologic findings. Contrary to our findings, a randomized crossover study found that a combination of MRCC and manual hyperinflation improved mucus clearance and oxygenation in subjects on mechanical ventilation.¹⁰ Manual hyperinflation, which involves lung ventilation by using a manual resuscitation bag, is a technique used in patients on mechanical ventilation to assist airway reopening.²¹ The reopening maneuver was not applied in our study, and MRCC alone could not remove the peripheral secretions nor re-expand the collapsed area. Therefore, a larger amount of mucus clearance might not lead to improved oxygenation.

MANUAL RIB CAGE COMPRESSION IN A PORCINE MODEL




Fig. 4. Changes in (A) atelectasis scores and (B) visual analog scale (VAS) scores in the study groups. Horizontal lines denote median value, the upper and lower edges represent the 25th and 75th percentiles, and whiskers represent the 10th and 90th percentiles. There were no significance differences between the 2 groups.

Table 2. Sequential Changes in Variables of Hemodynamics

Variable	After Mucus Infusion	0-1 Min	5 Min	10 Min	30 Min
Heart rate, beats/min					
Control group	129 ± 40	121 ± 37	123 ± 41	125 ± 41	123 ± 45
MRCC group	144 ± 26	128 ± 32	138 ± 30	141 ± 27	124 ± 18
Mean arterial pressure, mm Hg					
Control group	120 ± 14	115 ± 12	116 ± 11	118 ± 13	117 ± 12
MRCC group	124 ± 12	118 ± 19	127 ± 8	127 ± 9	122 ± 13
Cardiac output, L/min					
Control group	3.4 ± 0.8	3.4 ± 0.7	3.4 ± 0.7	3.3 ± 0.7	3.4 ± 0.8
MRCC group	3.3 ± 0.6	3.2 ± 0.5	3.2 ± 0.5	3.2 ± 0.6	3.4 ± 0.9

Data are shown mean ± standard deviation; there were no significant differences between groups. MRCC = manual rib cage compression

Although not significantly different between the groups, our results showed that MRCC exacerbated P_{aO_2} and P_{aCO_2} . A possible cause of this phenomenon is a decrease in end-expiratory lung volume. Furthermore, MRCC may exacerbate alveolar collapse (because of the compressive effect), which leads to decreasing the ventilation-perfusion ratio. In our study, all the pigs received mechanical ventilation with PEEP of 10 cm H_2O , and the protective effects of PEEP gradually stabilized the collapsed area; thus, P_{aO_2} increased in the MRCC group at the 30-min point. Another possible explanation is that an insufficient amount of mucus was removed from the airway due to the limitations of the suction protocol in our study. Increasing PEF contributes to moving mucus from the periphery to the proximal airways. Mucus retained in the proximal airway may influence ventilation and resistance parameters. Further studies are necessary to assess longer time courses or improved suctioning protocol to remove airway mucus completely.

We found no changes in hemodynamic parameters, in agreement with a previous study.¹² In general, vigorous physical stimuli such as MRCC can cause sympathetic activation and result in tachycardia and hypertension in conscious

patients. In the present study, however, the animals were anesthetized by using midazolam and ketamine, followed by a continuous infusion of pentobarbital and a muscle relaxant, vecuronium. Hence, it is likely that the pigs could hardly perceive physical stimulation associated with MRCC. Randomized trials have found that the heart rate was slightly but significantly reduced compared with controls in the post-manuever measurement.^{22,23} MRCC imposes external pressure on the rib cage and may change venous return. Nevertheless, the repercussion of the maneuver on the main hemodynamic variables was low. In addition, it is important to note that more significant hemodynamic changes may also occur in patients with hemodynamic instability.

Limitations

There were a few limitations of this study to consider. First, to assess the effects of MRCC on expiratory flow and avoid potential biases due to the animals' spontaneous exhalation, the pigs were paralyzed by using neuromuscular blockade. Thus, our findings are not entirely applicable to patients with spontaneous breathing. However, recent clinical studies investigated the effects of MRCC in specific conditions such as those that require a neuromuscular

MANUAL RIB CAGE COMPRESSION IN A PORCINE MODEL

blockade²⁴ or mucus hypersecretion.²⁵ Our experiment may contribute qualitative principles to consider the effectiveness of the intervention in specific conditions. Second, we were unable to assess atelectasis by using computed tomography due to a lack of equipment. Instead, we assessed atelectasis with visual analog scale scores from 0 to 10 cm by marking a single line on a scale anchored by the terms blackness and whiteness. Thus, the diagnosis of atelectasis may have lacked optimal sensitivity. Third, our study did not apply manual or ventilator hyperinflation; further research is needed to assess the effect of combining hyperinflation and MRCC. Fourth, our study demonstrated the safety of the intervention in terms of hemodynamic parameter; however, in elderly individuals who are debilitated or frail, we need to take into consideration the possibility of rib injury. Fifth, although pigs are commonly used to study airway mucus clearance, there seems to be considerable differences in morphology and function, such as collateral ventilation, among different species.²⁶ Therefore, translational clinical studies require further corroboration in humans.

Conclusions

Our findings indicated that hard MRCC combined with closed suctioning was safe and led to improvement of mucus clearance; however, no effectiveness was confirmed with regard to oxygenation and ventilation. Further well-conducted clinical studies are needed to thoroughly investigate the effects of MRCC.

ACKNOWLEDGMENTS

We thank Thomas Mayers, Medical English Communications Center, University of Tsukuba, for grammatical revision of this paper.

REFERENCES

- Konrad F, Schreiber T, Brecht-Kraus D, Georgieff M. Mucociliary transport in ICU patients. *Chest* 1994;105(1):237-241.
- Jelic S, Cunningham JA, Factor P. Clinical review: airway hygiene in the intensive care unit. *Crit Care* 2008;12(2):209.
- Hess DR. The evidence for secretion clearance techniques. *Respir Care* 2001;46(11):1276-1293.
- Hess DR. Airway clearance: physiology, pharmacology, techniques, and practice. *Respir Care* 2007;52(10):1392-1396.
- Tung A, Morgan SE. Modeling the effect of progressive endotracheal tube occlusion on tidal volume in pressure-control mode. *Anesth Analg* 2002;95(1):192-197, table of contents.
- Kawati R, Vimlati L, Guttman J, Hedenstierna G, Sjöstrand U, Schumann S, Lichtwarck-Aschoff M. Change in expiratory flow detects partial endotracheal tube obstruction in pressure-controlled ventilation. *Anesth Analg* 2006;103(3):650-657.
- Ntoumenopoulos G. Clinical impact of secretion retention. *Curr Respir Med Rev* 2014;10(3):158-162.
- Branson RD. Secretion management in the mechanically ventilated patient. *Respir Care* 2007;52(10):1328-1342; discussion 1342-1347.
- Unoki T, Kawasaki Y, Mizutani T, Fujino Y, Yanagisawa Y, Ishimatsu S, et al. Effects of expiratory rib-cage compression on oxygenation, ventilation, and airway-secretion removal in patients receiving mechanical ventilation. *Respir Care* 2005;50(11):1430-1437.
- Guimariães FS, Lopes AJ, Constantino SS, Lima JC, Canuto P, de Menezes SL. Expiratory rib cage compression in mechanically ventilated subjects: a randomized crossover trial [corrected]. *Respir Care* 2014;59(5):678-685.
- MacLean D, Drummond G, Macpherson C, McLaren G, Prescott R. Maximum expiratory airflow during chest physiotherapy on ventilated patients before and after the application of an abdominal binder. *Intensive Care Med* 1989;15(6):396-399.
- Marti JD, Li Bassi G, Rigol M, Saucedo L, Ranzani OT, Esperatti M, et al. Effects of manual rib cage compressions on expiratory flow and mucus clearance during mechanical ventilation. *Crit Care Med* 2013;41(3):850-856.
- Unoki T, Mizutani T, Toyooka H. Effects of expiratory rib cage compression combined with endotracheal suctioning on gas exchange in mechanically ventilated rabbits with induced atelectasis. *Respir Care* 2004;49(8):896-901.
- Unoki T, Mizutani T, Toyooka H. Effects of expiratory rib cage compression and/or prone position on oxygenation and ventilation in mechanically ventilated rabbits with induced atelectasis. *Respir Care* 2003;48(8):754-762.
- Comaru T, Fiori HH, Fiori RM, Padoim P, Stivanin JB, da Silva VD. Experimental model of atelectasis in newborn piglets. *Respir Care* 2014;59(1):121-126.
- American Association for Respiratory Care. AARC Clinical Practice Guidelines. Endotracheal suctioning of mechanically ventilated patients with artificial airways 2010. *Respir Care* 2010;55(6):758-764.
- Richter Larsen K, Ingwersen U, Thode S, Jakobsen S. Mask physiotherapy in patients after heart surgery: a controlled study. *Intensive Care Med* 1995;21(6):469-474.
- Benjamin RG, Chapman GA, Kim CS, Sackner MA. Removal of bronchial secretions by two-phase gas-liquid transport. *Chest* 1989;95(3):658-663.
- Kim CS, Rodriguez CR, Eldridge MA, Sackner MA. Criteria for mucus transport in the airways by two-phase gas-liquid flow mechanism. *J Appl Physiol* (1985) 1986;60(3):901-907.
- Volpe MS, Adams AB, Amato MB, Marini JJ. Ventilation patterns influence airway secretion movement. *Respir Care* 2008;53(10):1287-1294.
- Hodgson C, Denehy L, Ntoumenopoulos G, Santamaria J, Carroll S. An investigation of the early effects of manual lung hyperinflation in critically ill patients. *Anaesth Intensive Care* 2000;28(3):255-261.
- dos Santos RS, Donadio MV, da Silva GV, Blattner CN, Melo DA, Nunes FB, et al. Immediate effects of chest physiotherapy on hemodynamic, metabolic, and oxidative stress parameters in subjects with septic shock. *Respir Care* 2014;59(9):1398-1403.
- Bousari MP, Shirvani Y, Agha-Hassan-Kashani S, Nasab NM. The effect of expiratory rib cage compression before endotracheal suctioning on the vital signs in patients under mechanical ventilation. *Iran J Nurs Midwifery Res* 2014;19(3):285-289.
- Freynt A, Gobaille G, Joannes-Boyau O, Grandet P, Fleureau C, Ripoche J, et al. Effects of chest physiotherapy by expiratory flow increase on secretion removal and lung mechanics in ventilated patients: a randomized crossover study. *Intensive Care Med* 2016;42(6):1090-1091.
- Gonçalves EC, Souza HC, Tambascio J, Almeida MB, Basile Filho A, Gastaldi AC. Effects of chest compression on secretion removal, lung mechanics, and gas exchange in mechanically ventilated patients: a crossover, randomized study. *Intensive Care Med* 2016;42(2):295-296.
- Macklem PT. Airway obstruction and collateral ventilation. *Physiol Rev* 1971;51(2):368-436.