

Effects of 0 PEEP and $< 1.0 F_{IO_2}$ on S_{pO_2} and P_{ETCO_2} During Open Endotracheal Suctioning

Jacqueline Rodrigues de Freitas Vianna, Valéria Amorim Pires Di Lorenzo, Miléa Mara Lourenço da S Simões, Jorge Luís Guerra, and Maurício Jamami

BACKGROUND: Hyperoxygenation and hyperinflation, preferably with a mechanical ventilator, is the most commonly used technique to prevent the adverse effects of open endotracheal suctioning on arterial oxygenation and pulmonary volume. However, limited data are available on the effects of oxygen concentrations $< 100\%$ and PEEP with zero end-expiratory pressure (0 PEEP) to improve oxygenation and to maintain adequate ventilation during open endotracheal suctioning. The aim of this study was to analyze the behavior of S_{pO_2} and end-tidal CO_2 pressure (P_{ETCO_2}) in open endotracheal suctioning using the 0 PEEP technique with baseline F_{IO_2} (0 PEEP baseline F_{IO_2}) and 0 PEEP + hyperoxygenation of 20% above the baseline value (0 PEEP $F_{IO_2} + 0.20$) in critically ill subjects receiving mechanical ventilation. **METHODS:** This was a prospective, randomized, single-blind crossover study, for which 48 subjects with various clinical and surgical conditions were selected; of these, 38 subjects completed the study. The subjects were randomized for 2 interventions: 0 PEEP baseline F_{IO_2} and 0 PEEP $F_{IO_2} + 0.20$ during the open endotracheal suctioning procedure. Oxygenation was assessed via oxygen saturation as measured with pulse oximetry (S_{pO_2}), and changes in lung were monitored via P_{ETCO_2} using volumetric capnography. **RESULTS:** In the intragroup analysis with 0 PEEP baseline F_{IO_2} , there was no significant increase after open endotracheal suctioning in either S_{pO_2} ($P = .63$) or P_{ETCO_2} ($P = .11$). With 0 PEEP $F_{IO_2} + 0.20$, there was a significant increase in S_{pO_2} ($P < .001$), with no significant changes in P_{ETCO_2} ($P = .55$). In the intergroup comparisons, there was a significant increase compared to the basal values only with the 0 PEEP + 0.20 method at 1 min after hyperoxygenation ($P < .001$), post-immediately ($P < .001$), at 1 min after ($P < .001$), and at 2 min after open endotracheal suctioning ($P < .001$). **CONCLUSIONS:** The appropriate indication of the hyperinflation strategy via mechanical ventilation using 0 PEEP with or without hyperoxygenation proved to be efficient to maintain S_{pO_2} and P_{ETCO_2} levels. These results suggest that the technique can minimize the loss of lung volume due to open endotracheal suctioning. (ClinicalTrials.gov registration NCT02440919). *Key words:* suction; mechanical ventilation; pulmonary ventilation; physical therapy modalities; oxygen; capnography. [Respir Care 0;0(0):1–●. © 0 Daedalus Enterprises]

Introduction

Critically ill patients with artificial airways have shown changes in mucociliary clearance with the accumulation of secretions due to sedation, depression of the cough reflex,

high concentrations of oxygen, inadequate cuff pressure, inflammation, and mucosal trauma.¹⁻⁴ Airway aspiration is often necessary to maintain alveolar permeability and to prevent respiratory tract infection.⁵ According to the guidelines of the American Association for Respiratory Care (AARC),⁶ this procedure may result in numerous side effects, such as risks and complications for developing

Drs Rodrigues de Freitas Vianna, Pires Di Lorenzo, and Jamami are affiliated with the Department of Physical Therapy, Federal University of São Carlos, São Carlos, São Paulo, Brazil. Dr Rodrigues de Freitas Vianna, Ms Lourenço da S Simões, and Mr Guerra are affiliated with Intensive Care Physiotherapy, Santa Casa de Misericórdia Hospital, Batatais, São Paulo, Brazil.

Correspondence: Jacqueline Rodrigues de Freitas Vianna PT PhD, Av Comandante Salgado, 428 B, Castelo, Batatais, CEP 14300 220, São Paulo, Brazil. E-mail: jacrfvianna@uol.com.br.

The authors have disclosed no conflicts of interest.

DOI: 10.4187/respcare.07435

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

hypoxia/hypoxemia due to reduced level of tissue oxygenation (P_{aO_2}), decreased arterial oxygen saturation (S_{aO_2}), and peripheral oxygen saturation (S_{pO_2});⁷⁻¹¹ tissue trauma to the tracheal or bronchial mucosa with hemorrhagic secretion;⁷ increased microbial colonization of lower airways;⁹ decrease in dynamic compliance¹⁰⁻¹² and functional residual capacity;¹³ atelectasis;^{1,9} bronchoconstriction;¹⁴ changes in cerebral blood flow and increased intracranial pressure; vasovagal reactions; changes in blood pressure and heart rate;⁷⁻¹⁶ cardiac arrhythmias;¹⁵ low tidal volume;^{10,17-19} and elevated carbon dioxide levels in the blood (P_{aCO_2}).^{11,17}

Use of a closed suction system is suggested for adults with high F_{IO_2} or PEEP at risk of lung de-recruitment and for neonates.⁶ Advantages of a closed suction system compared with an open system include continuous mechanical ventilation, reduced hemodynamic impairment, shorter hospital stay, and reduced costs for patients and the health care system. Therefore, this method can replace open suction methods in the care of critically ill patients.²⁰

The recommendations to minimize or prevent a decrease in S_{aO_2} include appropriate catheter size, depth and time of suctioning, pressure setting, as well as routinely avoiding saline instillation and manual hyperinflation.^{8,17,18,20-23}

Hyperoxygenation is a method used to increase F_{IO_2} above basal levels, and 100% hyperoxygenation has been the most widely used method.^{6,16} Manual hyperinflation and ventilator hyperinflation are 2 physiotherapy procedures used to remove secretions. Manual hyperinflation with manual chest compression, also known as bag squeezing, was first described in 1968 by Clement and Hübsch.²⁴ This physiotherapeutic technique aims to improve oxygenation, clear bronchial secretions, and achieve alveolar re-expansion. Ventilator hyperinflation was originally described by Berney and Denehy²⁵ in 2002 as an alternative to manual lung hyperinflation, and it has been adopted to achieve the same purposes.

Based on the goal to increase expiratory air flow to remove secretions, studies have shown that pulmonary secretion removal depends not only on high expiratory flows, but also on the presence of an expiratory flow bias, ie, on the peak expiratory flow being higher than the peak inspiratory flow generated in the airways.²⁶ Furthermore, Volpe et al²⁷ reported that the most significant threshold for expiratory flow bias to move secretion toward the glottis, for human conditions, is the difference between peak expiratory flow and peak inspiratory flow being > 33 L/min.

The effects of 0 PEEP on lung compliance,^{4,28,30} gas exchange,^{4,28-30} tidal volume,³⁰ and hemodynamic repercussions^{4,28,29} have also been analyzed. This technique consists of imposing a gradual PEEP increase to 15 cm H₂O followed by an abrupt PEEP reduction to 0 PEEP in association with a manual bilateral thoracic compression to potentiate the increase of expiratory air flow, limiting the peak pressure to 40 cm H₂O.^{4,28-30} The ventilator hyperinflation with increased volume or with 0 PEEP was similar

QUICK LOOK**Current knowledge**

Open endotracheal suctioning is associated with hypoxemia and a loss of lung volume. Hyperinflation strategies using the mechanical ventilator with the 0 PEEP technique can maintain adequate levels of oxygen (S_{pO_2}) and minimize pulmonary ventilation loss (P_{ETCO_2}).

What this paper contributes to our knowledge

The 0 PEEP technique with or without hyperoxygenation maintained adequate S_{pO_2} and P_{ETCO_2} levels during open endotracheal suctioning. However, 0 PEEP with hyperoxygenation + 0.20 above baseline F_{IO_2} should be considered in subjects previously diagnosed with hypoxemia since it avoided the loss of $S_{pO_2} < 94\%$, even in subjects with $S_{pO_2} \leq 88\%$.

to manual hyperinflation in terms of the bronchial secretion removal, oxygenation, and with insignificant hemodynamic repercussions.^{4,28,29,31} The 0 PEEP technique appears to be safe, without alterations of hemodynamic variables, even in post-cardiac surgery patients.⁴ However, there have been few studies investigating using the 0 PEEP method to prevent hypoxemia or evaluating its impact on ventilation, whether associated with pre-oxygenation or not. Study designs involving ventilator hyperinflation are quite different, and the routine use of ventilator hyperinflation as well as the need for 100% hyperoxygenation are still debatable clinical issues.

Previous studies have reported that $F_{IO_2} = 1.0$ should be the method of choice to prevent lower levels of P_{aO_2} or S_{pO_2} , especially during open endotracheal suctioning, as recommended by the latest guidelines of the AARC.^{16,32,33} However, other studies suggest the evaluation of the need for hyperoxygenation with $F_{IO_2} = 1.0$, considering that the delivery of low or no oxygen was able to prevent hypoxemia during open endotracheal suctioning.^{11,34-39} Although O₂ was delivered for a short period of time, it is known that exposure to $F_{IO_2} = 1.0$ in humans produces toxic effects that can occur by reabsorption atelectasis, hyperoxic hypercapnia, bronchial and epithelial damage, decreased effectiveness of the ciliary epithelium, and bactericidal bronchial function.⁴⁰ Respiratory markers of oxidative stress were observed in healthy volunteers,⁴¹ while hyperoxygenation with 28% above baseline was delivered for 30 min.⁴¹ Oxidative damage was also detected in the lungs of rats exposed to 100% hyperoxygenation for 10 min.⁴²

Hyperoxygenation combined with hyperinflation has been the most commonly used technique for the prevention of the adverse effects caused by open endotracheal


PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

Fig. 1. Flow chart.

suctioning and is well documented in the literature.³² However little is known about the effects of < 100% oxygen delivery combined with the 0 PEEP technique on increasing oxygenation and maintaining adequate ventilation during open endotracheal suction. Therefore, we hypothesized that 0 PEEP baseline F_{IO_2} or 0 PEEP $F_{IO_2} + 0.20$ may maintain adequate S_{pO_2} and P_{ETCO_2} in open endotracheal suctioning without $F_{IO_2} = 1.0$ and the loss of lung volume. The aim of this study was to investigate the behavior of S_{pO_2} and P_{ETCO_2} in open endotracheal suctioning using 0 PEEP baseline F_{IO_2} and 0 PEEP $F_{IO_2} + 0.20$ in critically ill subjects on mechanical ventilatory support.

Methods

Study Location and Subject Population

The study was approved by the Human Research Ethics Committee of our institution and by the National Unified Research Registry involving human beings (Brazil Platform - CEP/Conep System- CAAE: 11354813.1.0000.5504). Written informed consent was obtained from the legal representatives of the subjects. The study was conducted in the tertiary adult ICU at Santa Casa de Misericórdia, Batatais, São Paulo, Brazil. All consecutive subjects requiring mechanical ventilation and ≥ 18 y of age were included during the period from June 2013 to May 2015. A total of 48 subjects of both sexes with several clinical and surgical conditions were selected. Of these, 10 subjects were excluded, leaving 38

subjects who completed the study. Figure 1 provides a flow chart of the study.

Inclusion and Exclusion Criteria

Inclusion criteria were as follows: subjects undergoing orotracheal intubation and on mechanical ventilation for > 12 h, hemodynamically stable, and requiring endotracheal suctioning according to the American Association for Respiratory Care criteria.⁶ Exclusion criteria were as follows: individuals using high doses of vasopressor amines or showing severe cardiac arrhythmias; with hemoglobin < 7 g/dL; impossibility of appropriate S_{pO_2} monitoring; baseline $F_{IO_2} \geq 0.60$; requirement of PEEP of > 10 cm H₂O; ARDS; rib fractures; presence of a chest drain; severe bronchospasm; intracranial hypertension (ie, intracranial pressure > 10 mm Hg); hemorrhagic disorders; marked degree of gastroesophageal reflux; bullous lung disease; unilateral lung disease; use of tracheostomy; closed suction system; peak pressure > 35 cm H₂O; hemodynamic instability with mean arterial pressure < 60 mm Hg; central venous pressure < 6 mm Hg; and no indication for endotracheal suctioning.

Study Design and Data Collection

This was a prospective, randomized, single-blind, cross-over study with randomization by drawing lots using opaque envelopes containing sequential interventions: Intervention I was 0 PEEP with F_{IO_2} (0 PEEP baseline


PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

Fig. 2. Data collection of experimental protocols. Baseline F_{IO_2} (current F_{IO_2} without hyperoxygenation); $F_{IO_2} + 0.20$ (hyperoxygenation with offer of F_{IO_2} 20% above baseline); Positioning (supine position for 20 min with the headrest at 40°); Pre (data collected at Baseline after positioning); Series: open endotracheal suctioning, data collected at 1 min after offer or not of oxygenation (Intervention I: 0 PEEP F_{IO_2} ; Intervention II: 0 PEEP $F_{IO_2} + 0.20$) and at 1 min before and after each catheter insertion, on the 5 open endotracheal suctioning procedures; After: data collected at 1 min, 2 min, and 30 min after open endotracheal suctioning.

F_{IO_2}), and Intervention II was 0 PEEP with 20% hyperoxygenation above baseline F_{IO_2} (0 PEEP $F_{IO_2} + 0.20$) during the open endotracheal suctioning procedure.

This study used the open suction system because it reflects the reality of Brazilian ICUs. The AARC guidelines were followed, not including adults with high F_{IO_2} and high PEEP levels with risk for lung de-recruitment.

To eliminate the carryover effects of the previous treatments, a washout period⁴³ of 5 h was allowed between interventions (Fig. 1). After the washout period, the subject was placed in a supine position for 20 min with the headrest at 40°. The endotracheal tube cuff was insufflated with 24 cm H₂O with a cuff pressure monitor (AMBU, Madrid, Spain). The following variables were then determined: tube size, level of consciousness, level of sedation with the vasoactive drug, indications for the open endotracheal suctioning procedure, ventilatory mode, ventilatory parameters, hemoglobin and hematocrit in the baseline condition before open endotracheal suctioning in both interventions. The interventions were performed by the senior author and a physiotherapist. Complications during or after the procedure, such as vomiting, trauma, or bronchospasms, were recorded.

S_{pO_2} was measured via pulse oximetry (Dixtal TECME S.A., Córdoba, Argentina) or ear lobe pulse oximetry (Dixtal TECME S.A., Córdoba, Argentina); the multiparameter monitor DX-2021 or DX-2023 (Dixtal TECME S.A.) was also used as an index of hypoxia. P_{ETCO_2} was measured via volumetric capnography with the Dixtal 3012 mechanical ventilator with a monitor and Capnostat 5 (Respironics,

Murrysville, Pennsylvania) and a sensor equipped with a mainstream flow sensor. Heart rate, breathing frequency, systolic arterial pressure, diastolic arterial pressure, and mean arterial pressure were obtained with the multiparameter monitor DX-2021 or DX-2023 (Capnostat 5, Respironics). S_{pO_2} and P_{ETCO_2} were measured at baseline (ie, immediately before each of the cleaning maneuvers) and then immediately after and again 30 min after the cleaning episodes were performed. S_{pO_2} was also measured during each of the cleaning episodes during both interventions at 1 min before oxygenation, 1 min after each of the 5 procedures for open endotracheal suctioning, and 1 min after oxygenation. At the end of data collection, subjects received the physiotherapy, medical, and nursing care routinely performed in the ICU (Fig. 2).

Procedures

All subjects were ventilated using the following ventilatory modes: volume control ventilation, assist/control; pressure control ventilation, assist/control; volume control synchronized intermittent mandatory ventilation; and spontaneous breath with volume guarantee pressure-support ventilation. After all inclusion criteria were checked, the subjects were randomly allocated to one of the 2 sequences. The subjects were placed in the supine position with elevation of the head of the bed at 40° for 20 min before the baseline measurements and the open endotracheal suctioning procedure.

The open endotracheal suction procedure was performed using an atoxic, sterile, siliconized polyvinyl chloride, 12

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

French catheter with a lateral and bottom orifice, and a suction control valve (Embramed, São Paulo- SP- Brazil) for endotracheal tubes with an internal diameter of 7.0–8.5 mm.⁴⁴ The catheter was introduced until resistance was met, and it was then withdrawn 2–3 cm with a negative pressure of ~150 mm Hg^{45,46} applied for 15 s with circular movements; there was a 60-s interval between each of the 5 insertions, according to the recommendations of the latest endotracheal suctioning directives of the American Association for Respiratory Care.⁶

Hyperoxygenation was performed using a Dixtal 3012 ventilator. In 0 PEEP hyperoxygenation, F_{IO₂} + 0.20 was increased to a value 20% above the baseline value of the subject for 1 min before each of the 5 procedures and for 1 min after the open endotracheal suctioning procedure.

At the start of each maneuver, PEEP was increased to 15 cm H₂O for 60 s at the end of the inspiratory phase, with the pressure peak limited to 40 cm H₂O.²⁵ PEEP was then abruptly reduced to 0 cm H₂O; this cycle is known as 0 PEEP.^{28–30} The tidal volume was previously adjusted to 6 mL/kg with an inspiratory time of 0.8–1.2 s and a breathing frequency of 8 breaths/min, followed by an abrupt PEEP reduction to 0 cm H₂O (0 PEEP) in the expiratory phase, returning the PEEP value to the inspiratory phase. The maneuver was repeated after 2 ventilatory cycles at baseline PEEP. In each step, the maneuver was performed 3 times^{28–30,47,48} before each of the 5 open endotracheal suctioning procedures, for a total of 15 cycles per step, totaling 15 min.

Statistical Analysis

The sample size calculation was performed using GPower 3.1 with the dependent variables (S_{pO₂} and P_{ETCO₂}) of a pilot study with 16 subjects carried out by the authors, where $\alpha = 5\%$ (2-tailed) and 80% power, estimating a sample of 41 subjects. Data were analyzed using GraphPad Prism 5.01 (GraphPad Software, San Diego, California). Initially, 1-way repeated measures analysis of variance was used to test the possibility of grouping all subjects within different ventilation modes. According to the Shapiro-Wilk test for sample data with non-normal distribution, we used the paired Wilcoxon test for between and within comparisons. The Friedman test post hoc Dunn test was used for the 4 measures obtained during the 5 open endotracheal suctioning procedures. A significance level was set at 5% for all tests. Data are expressed as absolute frequency (*n*), percentage (%), or median (range).

Results

Thirty-eight subjects with mean age of 65.82 ± 12.26 y participated in the study. The initial sample characteristics are displayed in Table 1. The indication for mechanical

Table 1. Subject Characteristics and Baseline Conditions

Variable	<i>n</i> (%)
Sex	
Female	12 (31.6)
Male	26 (68.4)
Causes of intubation	
Respiratory	24 (63)
Cardiac	11 (29)
Surgical (abdominal and orthopedic)	3 (8)
Ventilation mode	
Volume control ventilation, assist/control	10 (26.3)
Pressure control ventilation, assist/control	5 (13.2)
Volume control synchronized intermittent mandatory ventilation	8 (21.1)
Pressure control synchronized intermittent mandatory ventilation	8 (21.1)
Volume-guaranteed pressure support	7 (18.4)

N = 38 subjects.

ventilation was predominantly respiratory, mainly due to COPD and acute respiratory failure caused by pneumonia.

The Acute Physiologic and Chronic Health Evaluation II (APACHE II) prognostic index was applied upon admission to the hospital and 48 h after admission to estimate the severity of the disease and predict hospital mortality. The mean APACHE II scores were 22.2 ± 8.64 and 22.3 ± 8.28. The risk of death was 40% for nonoperative subjects and 30% for postoperative subjects. The mortality rate was 55.3% (*n* = 21), with a high rate of ICU discharge of 44.7% (*n* = 17). The mean duration of mechanical ventilation was 14.29 ± 12.81 d, and the mean length of stay in the ICU was 14.72 ± 9.08 d.

The endotracheal tubes had a mean internal diameter size of 7.99 ± 0.36 mm (range 7.0–8.5 mm). Most tubes (66%) used were 8.5 mm in diameter. There was no statistically significant difference between F_{IO₂}, PEEP, mean airway pressure, and hemoglobin and hematocrit values in baseline conditions (*P* > .05) (Table 2).

Oxygenation Behavior Measured With S_{pO₂}

There was a significant increase in S_{pO₂} within groups immediately after the open endotracheal suctioning with the 0 PEEP F_{IO₂} + 0.20 method, with S_{pO₂} returning to baseline values after 30 min in both interventions; however, the S_{pO₂} values were higher with the 0 PEEP F_{IO₂} + 0.20 method compared to the 0 PEEP baseline F_{IO₂} method (Table 3). In the continuous analyses within groups, there was a significant increase compared to the basal values only in the 0 PEEP F_{IO₂} + 0.20 method at 1 min after hyperoxygenation (*P* < .001), post-immediately (*P* < .001), at 1 min after (*P* < .001), and at 2 min after open endotracheal suctioning (*P* < .001) (Fig. 3).

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

Table 2. Comparison of Subjects' Basal Conditions

	Intervention I	Intervention II	P*
Hemoglobin, g/dL	10.9 (7.2–15.4)	11.5 (7.0–15.2)	.26
Hematocrit, %	34.7 (23.5–53.5)	35.9 (21.0–53.7)	.84

Data are presented as median (range). N = 38 subjects. Intervention I: 0 PEEP, baseline F_{IO₂}; Intervention II: 0 PEEP, hyperoxygenation.
*Paired Wilcoxon test used for statistical comparison.

In both interventions, the 4 values were maintained during the 5 suction procedures ($P = .65$ and $P = .93$). A significant difference was found between the groups after hyperoxygenation ($P = .003$) and at all time points measured ($P < .001$). No significant difference was observed in baseline values and at 30 min after the procedure (Fig. 3).

It is noteworthy that, in the 0 PEEP baseline F_{IO₂} intervention, 6 subjects (15.8%) had hypoxemia ($S_{pO_2} < 90\%$), 2 subjects had $S_{pO_2} \leq 90\%$ during and after 2 min, and 3 (7.8%) subjects had $S_{pO_2} \leq 90\%$ after 30 min of open endotracheal suctioning. In the 0 PEEP F_{IO₂} + 0.20 method, 4 subjects had baseline hypoxemia (10.5%) ($S_{pO_2} < 90\%$) and 1 subject (7.8%) exhibited $S_{pO_2} = 90\%$ after 30 min following open endotracheal suctioning.

Ventilation Behavior Evaluated With P_{ETCO₂}

In the intragroup analysis, no significant changes were observed in P_{ETCO₂} at any time point in either intervention ie. ($P = .11$, $P = .46$, $P = .29$, and $P = .32$); the same was noted in the between-group analysis ie. ($P = .10$, $P = .50$ and $P = .98$) (Table 3, Fig. 4). Emphasis should be given to the baseline homogeneity of the interventions. In 0 PEEP baseline F_{IO₂}, 27 subjects (71%) were isocapnic (P_{ETCO₂} ≤ 45 mm Hg) and 11 (29%) were hypercapnic (P_{ETCO₂} > 45 mm Hg); in 0 PEEP F_{IO₂} + 0.20, 26 subjects were isocapnic (68%) and 12 were hypercapnic (32%).




Fig. 3. Changes in S_{pO_2} before and after open endotracheal suctioning at different time points with 0 PEEP F_{IO₂} + 0.20 and 0 PEEP baseline F_{IO₂}. †Comparison between the groups. ‡Comparison within the groups pre- and immediately after.

Discussion

In this study, we analyzed the behavior of S_{pO_2} and P_{ETCO₂} in open endotracheal suctioning using the 0 PEEP technique with F_{IO₂} at baseline and 0 PEEP + hyperoxygenation of 20% above the baseline value in critically ill subjects receiving mechanical ventilation. The results indicated that the 0 PEEP technique with or without hyperoxygenation during the open endotracheal suctioning procedure was effective in maintaining adequate oxygenation, with a median S_{pO_2} of 95%, as well as adequate ventilation, with a median P_{ETCO₂} of 39.5 mm Hg. The PEEP to 0 PEEP F_{IO₂} + 0.20 technique maintained adequate S_{pO_2} even in subjects previously diagnosed with hypoxemia ($S_{pO_2} \leq 88\%$).

S_{pO_2} is one of the most important advances in respiratory monitoring. It is commonly used to assess oxygenation of ICU subjects during open endotracheal suctioning, mainly

Table 3. Changes in S_{pO_2} and P_{ETCO₂} During the Open Endotracheal Suctioning Procedure

Tempo	S_{pO_2} , %			P _{ETCO₂} , mm Hg		
	Intervention I	Intervention II	P*	Intervention I	Intervention II	P*
Pre	0.94 (0.88–0.99)	0.95 (0.87–1.0)	.19	39.5 (15–72)	40.5 (16–68)	.10
Immediately after	0.95 (0.87–1.0)	0.99 (0.93–1.0)	< .001	37 (11–75)	39.5 (16–71)	.050
Pre × Immediately after	P^\dagger	.63	< .001	.11	.29	
30 min after	0.96 (0.89–1.0)	0.95 (0.90–0.99)	.77	39 (12–63)	40 (17–66)	.98
Pre × 30 min after	P^\ddagger	.055	.89	.46	.32	

Data are presented as median (range). Intervention I: 0 PEEP, baseline F_{IO₂}; Intervention II: 0 PEEP, hyperoxygenation.

* Paired Wilcoxon test used for between comparison.

† Paired Student *t* test for within comparison (Pre × Immediately after).

‡ Paired Student *t* test for within comparison (Pre × 30 min after).


PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

Fig. 4. Changes in P_{ETCO₂} before and after open endotracheal suctioning at different time points with 0 PEEP F_{IO₂} + 0.20 and 0 PEEP baseline F_{IO₂}. †Comparison between the groups. ‡Comparison within the groups pre- and immediately after.

due to its the degree of accuracy and the ease of operation of most pulse oximetry instruments.^{10,50,51} The S_{pO₂} value usually reflects S_{aO₂}, especially when > 90%, with normal values of S_{pO₂} and S_{aO₂} ranging from 3% to 5%, respectively.^{50,52}

Helayel et al⁵³ reported that S_{pO₂} values ≥ 99% ruled out the possibility of S_{aO₂} < 90% in critically ill or surgical patients. Jubran et al⁵² considered that limits between 92% and 94% should be targets to avoid S_{aO₂} < 90%, which corresponds to mild hypoxemia.^{51,54} However, Seguin et al⁵⁵ detected the absence of hypoxemia only with S_{pO₂} values ≥ 96% in critically ill subjects receiving mechanical ventilation, ie, the value that would determine S_{pO₂} ≥ 90%. According to Torsani et al,⁵⁶ 88–92% S_{pO₂} is typically accepted as ideal in clinical practice for critically ill subjects. Nonetheless, Jubran⁵¹ reported that S_{pO₂} > 92% would be a satisfactory level of oxygenation for mechanically ventilated subjects. Ghayumi et al⁵⁷ recently reported that S_{pO₂} ≥ 94% can be used as a reliable and accurate substitute for arterial blood gas for the assessment of hypoxemia and that arterial blood gas analysis should be limited to subjects with S_{pO₂} < 94%.

In this study, when considering the moments before and immediately after open endotracheal suctioning, the median S_{pO₂} in 0 PEEP baseline F_{IO₂} was 94–95%; with the 0 PEEP + hyperoxygenation of 20% method, it was 95–99%, maintaining oxygenation with or without hyperoxygenation. It should be noted that the 0 PEEP baseline F_{IO₂} method was effective in preventing levels of S_{pO₂} ≥ 97%.

However, in the 0 PEEP baseline F_{IO₂} method, 4 subjects (10%) presented with baseline S_{pO₂} = 88% before open endotracheal suctioning, and 3 subjects maintained S_{pO₂} ≤

90% during the procedure. In the 0 PEEP hyperoxygenation method, 3 subjects (8%) exhibited baseline S_{pO₂} ≤ 88%. S_{pO₂} ≥ 94% was maintained in all subjects during the procedure.

Clinical trials have indicated that there are no beneficial effects of hyperoxia on critically ill subjects.⁴⁰ Prescription of F_{IO₂} levels higher than are necessary to achieve arterial oxygenation goals may further increase the risk of oxygen toxicity.⁵¹ The use of F_{IO₂} = 1.0 has potential toxic effects due to the release of large amounts of proinflammatory cytokines and significant microvascular and muscular changes, inducing severe lung injury.^{2,40,42,58-60}

We investigated the delivery of 20% above baseline F_{IO₂}, similar to a previous reported by Rogge et al³⁴ in 11 subjects with COPD, who compared the delivery of 20% above baseline F_{IO₂} to the delivery of F_{IO₂} = 1.0 both combined with manual lung hyperinflation, observing whether the O₂ levels were properly maintained in both oxygen therapies.

Souza et al³⁷ reported the effectiveness of F_{IO₂} + 0.20 delivery by comparing baseline F_{IO₂} versus hyperoxygenation with F_{IO₂} + 0.20 above baseline, combined with conventional lung physiotherapy and bag squeezing technique in 30 critically ill subjects. Demir and Dramali³⁵ also reported that S_{pO₂} and P_{aO₂} without hyperoxygenation was not significantly reduced in intubated subjects undergoing mechanical ventilation with closed endotracheal suctioning.

In the present study, the 0 PEEP technique with baseline F_{IO₂} and the 0 PEEP F_{IO₂} + 0.20 technique in open endotracheal suctioning maintained adequate levels of S_{pO₂} in critically ill subjects receiving mechanical ventilation.

Capnography is an effective noninvasive method in diagnosing early respiratory depression and airway disorders. It also indirectly measures metabolism and circulation, estimating P_{aCO₂} with some accuracy, whereas the observed gradient is minimal, with a significant difference of 2–5 mm Hg when comparing P_{aCO₂} and P_{ETCO₂}. A normal average P_{aCO₂} value of 40 mm Hg has a capnometry reading of 35–38 mm Hg P_{ETCO₂}.⁶¹⁻⁶⁶

Savian et al³¹ observed an increase in CO₂ production with the application of manual lung hyperinflation in subjects under the volume control synchronized intermittent mandatory ventilation mode. Ahmed et al⁶⁷ also reported that ventilator hyperinflation provided better dynamic compliance compared to manual hyperinflation.

Our study indicated that both interventions (ie, 0 PEEP baseline F_{IO₂} and 0 PEEP F_{IO₂} + 0.20) maintained adequate levels of P_{ETCO₂} in critically ill subjects undergoing mechanical ventilation in the moments before and after open endotracheal suctioning, suggesting that the 0 PEEP technique can minimize the loss of lung volume due to pulmonary de-recruitment with an open endotracheal suctioning system.

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

Berney and Denehy²⁵ and Savian et al³¹ described that both hyperinflation methods (ie, manual hyperinflation and ventilator hyperinflation) proved to be effective in removing secretions and improving static lung compliance, hemodynamics, and oxygenation. We used the 0 PEEP technique, which was applied in only 3 studies without hyperoxygenation, following the same patterns; ie, imposing a gradual PEEP increase to 15 cm H₂O, with peak pressure limited to 40 cm H₂O for 5 ventilatory cycles, followed by an abrupt PEEP reduction. The variations found were the different evaluation time periods (5 min, 10 min, or 3 sets of 5 ventilation cycles) with or without manual chest compression.

Rodrigues²⁸ reported that the 0 PEEP technique in combination with manual chest compression was a safe technique in removing bronchial secretions in myocardial revascularization subjects during the immediate postoperative period. The 0 PEEP technique also allowed greater control in respiratory mechanics, monitoring pressures and flows delivered to the lungs. Santos et al³⁰, using the same study design²⁸ to compare the 0 PEEP technique combined with manual chest compressions in 12 subjects undergoing mechanical ventilation, reported a significant increase in the respiratory system compliance in both techniques, with no significant differences between them as well as favorable behavior of S_{pO₂} in the manual chest compressions group. In a similar study with 20 subjects, Lobo et al²⁹ compared the 0 PEEP technique combined with manual vibro-compression with no hyperoxygenation versus a bag-squeezing technique, using 5 L/m of oxygen (F_{IO₂} = ~0.40). The authors considered the 0 PEEP technique feasible in critically ill subjects because its use did not cause any significant hemodynamic repercussions. In 2011, Herbst-Rodrigues et al⁴ analyzed 15 subjects submitted to a coronary artery bypass graft surgery and did not find alterations in the hemodynamic variables (ie, S_{pO₂} and P_{ETCO₂}), confirming the safety of the 0 PEEP technique combined with manual chest compressions when it is not used as an alternative technique for the removal of bronchial secretions.

In the present study, the removal of bronchial secretions was not analyzed. We used the method of previous studies, without association with manual chest compression and with pressurization time and using 60 s in each series, and we noted similar results in terms for the alterations in S_{pO₂} and P_{ETCO₂}. The 0 PEEP technique in the expiratory flow bias concept is effective for mucus mobilization in the central direction, maintaining the levels of S_{pO₂} and P_{ETCO₂}.

It is noteworthy that the standardized 0 PEEP technique appears to be a safe method in cardiovascular subjects, given that no hemodynamic alterations were observed in those subjects in previous studies.^{4,28} The 0 PEEP application time should not exceed 15 min, because there could be repercussions for heart rate.⁴⁹ This technique should not be

used with subjects with COPD because potential detrimental effects of manual hyperinflation have been reported, including increased intrinsic PEEP and its consequences.⁶⁸ Further research focusing on the effects of the 0 PEEP method on subjects with COPD is required for a better understanding of the potential benefits of this technique.

Open endotracheal suctioning may cause a significant but transient loss of lung volume.⁶⁹ The prevention of lung volume loss is of paramount importance because the high shear forces between the open and closed lung units, combined with lung de-recruitment, can be harmful to the lungs.⁷⁰ Therefore, future studies should focus on the fact that lung volume loss should be avoided not only during but also after open endotracheal suctioning using lung recruitment strategies.

This study has limitations that are inherent to its design, although no significant differences were noted in baseline interventions and in the type of sample. A study population with more homogeneous clinical features is likely to provide a better understanding of the results. Additional investigations are needed to compare different ventilatory modes and strategies for the protection of pulmonary volume during and after open circuit suctioning, as well as the impact of the relationship between the diameter of the suction catheter and the endotracheal tube volume losses during open endotracheal suctioning.

Conclusions

The results indicated that an appropriate indication of the hyperinflation strategy by mechanical ventilation using the 0 PEEP technique with or without hyperoxygenation was efficient to maintain S_{pO₂} and P_{ETCO₂} levels in critically ill subjects, suggesting that the technique minimized lung volume loss due to the open endotracheal suctioning system. However, we recommend the use of the 0 PEEP technique + 20% above the baseline in subjects previously diagnosed with hypoxemia because the 0 PEEP with hyperoxygenation + 0.20 above baseline F_{IO₂} method avoided the loss of S_{pO₂} < 94%, even in subjects with S_{pO₂} ≤ 88%.

ACKNOWLEDGMENTS

The authors acknowledge the entire ICU team from Santa Casa de Batatais and their families. Thanks are also due to Érica Moreira Oliveira for her contribution to the pilot project.

REFERENCES

1. Sackner MA. Letter: tracheobronchial suction catheters. *Chest* 1976;70(2):318-320.
2. Landa JF, Kwoka MA, Chapman GA, Brito M, Sackner MA. Effects of suctioning on mucociliary transport. *Chest* 1980;77(2):202-207.

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

3. Keller C, Brimacombe J. Bronchial mucus transport velocity in paralyzed anesthetized patients: a comparison of the laryngeal mask airway and cuffed tracheal tube. *Anesth Analg* 1998;86(6):1280-1282.
4. Herbst-Rodrigues MV, Vo C, Auler JOC Jr, Feltrim M. PEEP-ZEEP technique: cardiorespiratory repercussions in mechanically ventilated patients submitted to a coronary artery bypass graft surgery. *J Cardiac Surgery* 2011;13(6):108-114.
5. Jelic S, Cunningham JA, Factor P. Clinical review: airway hygiene in the intensive care unit. *Crit Care* 2008;12(2):209-218.
6. American Association for Respiratory Care. AARC Clinical Practice Guidelines: endotracheal suctioning of mechanically ventilated patients with artificial airways 2010. *Respir Care* 2010;55(6):758-764.
7. Maggiore SM, Lellouche F, Pignataro C, Girou E, Maitre B, Richard JC, et al. Decreasing the adverse effects of endotracheal suctioning during mechanical ventilation by changing practice. *Respir Care* 2013;58(10):1588-1597.
8. Maggiore SM, Lellouche F, Pigeot J, Taille S, Deye N, Durrmeyer X, et al. Prevention of endotracheal suctioning-induced alveolar derecruitment in acute lung injury. *Am J Respir Crit Care Med* 2003;167(9):1215-1224.
9. Tingay DG, Copnell B, Grant CA, Dargaville PA, Dunster KR, Schibler A. The effect of endotracheal suction on regional tidal ventilation and end-expiratory lung volume. *Intensive Care Med* 2010;36(5):888-896.
10. Morrow B, Futter M, Argent A. Effect of endotracheal suction on lung dynamics in mechanically-ventilated paediatric patients. *Australian J Physiother* 2006;52(2):121-126.
11. Avena MJ, de Carvalho WB, Beppu OS. Evaluation of oxygenation, ventilation and respiratory mechanics before and after endotracheal suction in mechanically ventilated children. *Rev Assoc Med Bras (1992)* 2003;49(2):156-161.
12. Main E, Castle R, Newham D, Stocks J. Respiratory physiotherapy vs. suction: the effects on respiratory function in ventilated infants and children. *Intensive Care Med* 2004;30(6):1144-1151.
13. Heinze H, Sedemund-Adib B, Heringlake M, Gosch UW, Eichler W. Functional residual capacity changes after different endotracheal suctioning methods. *Anesth Analg* 2008;107(3):941-944.
14. Guglielminotti J, Desmots JM, Dureuil B. Effects of tracheal suctioning on respiratory resistances in mechanically ventilated patients. *Chest* 1998;113(5):1335-1338.
15. Seymour CW, Cross BJ, Cooke CR, Gallop RL, Fuchs BD. Physiologic impact of closed-system endotracheal suctioning in spontaneously breathing patients receiving mechanical ventilation. *Respir Care* 2009;54(3):367-374.
16. Bourgault AM, Brown CA, Hains SM, Parlow JL. Effects of endotracheal tube suctioning on arterial oxygen tension and heart rate variability. *Biol Res Nurs* 2006;7(4):268-278.
17. Lasocki S, Lu Q, Sartorius A, Fouillat D, Remerand F, Rouby JJ. Open and closed-circuit endotracheal suctioning in acute lung injury: efficiency and effects on gas exchange. *Anesthesiology* 2006;104(1):39-47.
18. Sakuramoto H, Shimojo N, Jesmin S, Unoki T, Kamiyama J, Oki M, et al. Repeated open endotracheal suctioning causes gradual desaturation but does not exacerbate lung injury compared to closed endotracheal suctioning in a rabbit model of ARDS. *BMC Anesth* 2013;13(1):47-57.
19. Briassoulis G, Briassoulis P, Michaeloudi E, Fitrolaki DM, Spanaki AM, Briassoulis E. The effects of endotracheal suctioning on the accuracy of oxygen consumption and carbon dioxide production measurements and pulmonary mechanics calculated by a compact metabolic monitor. *Anesth Analg* 2009;109(3):873-879.
20. Afshari A, Safari M, Oshvandi K, Soltanian AR. The effect of the open and closed system suction on cardiopulmonary parameters: time and costs in patients under mechanical ventilation. *Nurs Midwifery Stud* 2014;3(2):e14097.
21. Akgül S, Akyolcu N. Effects of normal saline on endotracheal suctioning. *J Clin Nurs* 2002;11(6):826-830.
22. Stiller K. Physiotherapy in intensive care: towards an evidence-based practice. *Chest* 2000;118(6):1801-1813.
23. Zeitoun SS, Barros A, Diccini S. A prospective, randomized study of ventilator-associated pneumonia in patients using a closed vs. open suction system. *J Clin Nurs* 2003;12(4):484-489.
24. Clement AJ, Hübsch SK. Chest physiotherapy by the 'bag squeezing' method: a guide to technique. *Physiotherapy* 1968;54(10):355-359.
25. Berney S, Denehy L. A comparison of the effects of manual and ventilator hyperinflation on static lung compliance and sputum production in intubated and ventilated intensive care patients. *Physiother Res Int* 2002;7(2):100-108.
26. Benjamin RG, Chapman GA, Kim CS, Sackner MA. Removal of bronchial secretions by two-phase gas-liquid transport. *Chest* 1989;95(3):658-663.
27. Volpe MS, Adams AB, Amato MB, Marini JJ. Ventilation patterns influence airway secretion movement. *Respir Care* 2008;53(10):1287-1294.
28. Rodrigues M. Estudo do comportamento hemodinâmico, da troca gasosa, da mecânica respiratória e da análise do muco brônquico na aplicação de técnicas de remoção de secreção brônquica em pacientes sob ventilação mecânica [Thesis]. São Paulo. Faculdade de Medicina da Universidade de São Paulo 2007.
29. Lobo DML, Cavalcante LA, Mont'Alverne DGB. Applicability of bag squeezing and zEEP maneuvers in mechanically ventilated patients. *Rev Bras Ter Intensiva* 2010;22(2):186-191.
30. Santos FRA, Schneider Junior LC, Forgiarini Junior LA, Veronezi J. Efeitos da compressão torácica manual versus a manobra de PEEP-ZEEP na complacência do sistema respiratório e na oxigenação de pacientes submetidos à ventilação mecânica invasiva. *Rev Bras Ter Intensiva* 2009;21(2):155-161.
31. Savian C, Paratz J, Davies A. Comparison of the effectiveness of manual and ventilator hyperinflation at different levels of positive end-expiratory pressure in artificially ventilated and intubated intensive care patients. *Heart Lung* 2006;35(5):334-341.
32. Oh H, Seo W. A meta-analysis of the effects of various interventions in preventing endotracheal suction-induced hypoxemia. *J Clin Nurs* 2003;12(6):912-924.
33. Jerre G, Beraldo MA, Silva T. D J, Gastaldi A, Kondo C, Leme F, et al. Fisioterapia no paciente sob ventilação mecânica. *Rev Bras Ter Intensiva* 2007;19(3):399-407.
34. Rogge JA, Bunde L, Baun MM. Effectiveness of oxygen concentrations of less than 100% before and after endotracheal suction in patients with chronic obstructive pulmonary disease. *Heart Lung* 1989;18(1):64-71.
35. Demir F, Dramali A. Requirement for 100% oxygen before and after closed suction. *J Adv Nurs* 2005;51(3):245-251.
36. Diniz G, Oliveira HMD, Arrais RCS, Pessoa BP, Figueiredo PHS. A utilização de FiO₂ inferior a 100% para hiperoxigenação de pacientes estáveis submetidos à aspiração endotraqueal. *ASSOBRAFIR Ciência* 2012;3(1):45-56.
37. Souza T, Souza R, Milani CM. Comportamento das variáveis cardiorrespiratórias de pacientes em ventilação mecânica submetidos à aspiração traqueal sob diferentes níveis de fração inspirada de oxigênio. *Revista Eletrônica Saúde e Ciência* 2012;2(5):63-77.
38. Overend TJ, Anderson CM, Brooks D, Cicutto L, Keim M, McAuslan D, Nonoyama M. Updating the evidence-base for suctioning adult patients: a systematic review. *Can Respir J* 2009;16(3):e6-e17.
39. Ntoumenopoulos G. Endotracheal suctioning may or may not have an impact, but it does depend on what you measure. *Respir Care* 2013;58(10):1707-1710.

PEEP AND O₂ DURING OPEN ENDOTRACHEAL SUCTIONING

40. Gordo-Vidal F, Calvo-Herranz E, Abella-Alvarez A, Salinas-Gabina I. Hyperoxia-induced pulmonary toxicity. *Med Int* 2010;34(2):134-138.
41. Phillips M, Cataneo RN, Greenberg J, Grodman R, Gunawardena R, Naidu A. Effect of oxygen on breath markers of oxidative stress. *Eur Respir J* 2003;21(1):48-51.
42. Valença S, Kloss ML, Bezerra FS, Lanzetti M, Silva FL, Porto LC. Efeitos da hiperóxia sobre o pulmão de ratos Wistar. *J Bras Pneumol* 2007;33(6):655-662.
43. Dawson-Saunders B, Trapp RG. Basic and clinical biostatistics, 2nd ed. Norwalk, CT: Appleton and Lange; 1994:14-44.
44. Liu XW, Jin Y, Ma T, Qu B, Liu Z. Differential effects of endotracheal suctioning on gas exchanges in patients with acute respiratory failure under pressure-controlled and volume-controlled ventilation. *Biomed Res Int* 2015;2015:941081.
45. Yousefi HVJ, Yazdannik AR. Comparison of the effects of two levels of negative pressure in open endotracheal tube suction on the physiological indices among patients in intensive care units. *Iran J Nurs and Midwifery Res* 2014;19(5):473-477.
46. Fernández MD, Piacentini E, Blanch L, Fernández R. Changes in lung volume with three systems of endotracheal suctioning with and without pre-oxygenation in patients with mild-to-moderate lung failure. *Intensive Care Med* 2004;30(12):2210-2215.
47. Chicayban LM, de Azevedo Silva M, Fagundes VB. PEEP-ZEEP com compressão torácica e hiperinsuflação com ventilador mecânico: ensaio clínico cruzado randomizado. *Rev Bras Fisioter* 2010;14(Suppl 1):529.
48. Auler JO Jr, Carmona MJ, Barbas CV, Saldiva PH, Malbouisson LM. The effects of positive end-expiratory pressure on respiratory system mechanics and hemodynamics in postoperative cardiac surgery patients. *Braz J Med Biol Res* 2000;33(1):31-42.
49. Auler JOC Jr, Franchi AA, Coimbra VRM, de Carvalho MJ. Estudo comparativo entre CO₂ obtido por capnografia e gasometria no pós-operatório de cirurgia cardíaca. *Rev Bras Anestesiol* 1990;40(5):311-318.
50. Coté CJ, Goldstein EA, Fuchsmann WH, Hoaglin DC. The effect of nail polish on pulse oximetry. *Anesth Analg* 1988;67(7):683-686.
51. Jubran A. Pulse oximetry. *Crit Care* 1999;3(2):R11-17.
52. Jubran A. Advances in respiratory monitoring during mechanical ventilation. *Chest* 1999;116(5):1416-1425.
53. Helayel PE, de Oliveira Filho GR, Marcon L, Pedemeiras FH, Nicolodi MA, Pedemeiras SG. Gradiente SpO₂ - SaO₂ durante ventilação mecânica em anestesia e terapia intensiva. *Rev Bras Anestesiol* 2001;51(4):305-310.
54. Jubran A, Tobin MJ. Reliability of pulse oximetry in titrating supplemental oxygen therapy in ventilator-dependent patients. *Chest* 1990;97(6):1420-1425.
55. Seguin P, Le Rouzo A, Tanguy M, Guillou YM, Feuillu A, Malledant Y. Evidence for the need of bedside accuracy of pulse oximetry in an intensive care unit. *Crit Care Med* 2000;28(3):703-706.
56. Torsani VMJ, Costa E. Monitorização na insuficiência respiratória aguda. *Pneumologia Paulista* 2013;27(1):31-33.
57. Ghayumi SM, Khalafi-Nezhad A, Jowkar Z. Pulse oximeter oxygen saturation in prediction of arterial oxygen saturation in liver transplant candidates. *Hepat Mon* 2014;14(3):e15449.
58. Lopes CR, Sales ALM, Simões MDJ, Angelis MA, Oliveira NML. Efeitos agudos da ventilação mecânica com hiperoxia na morfometria do diafragma de ratos. *Rev Bras Fisioter* 2009;13(6):487-492.
59. Pereira DS, Silva BAK, Aydos RD, Carvalho PTC, Odashiro AN, Oliveira L. Efeitos da alta concentração de oxigênio (hiperóxia) por tempo prolongado no tecido pulmonar de ratos wistar revista biociências. *UNITAU* 2008;14(2):100-116.
60. Bhandari V, Choo-Wing R, Lee CG, Zhu Z, Nedrelew JH, Chupp GL, et al. Hyperoxia causes angiotensin 2-mediated acute lung injury and necrotic cell death. *Nat Med* 2006;12(11):1286-1293.
61. McArthur CD, AARC. AARC Clinical Practice Guideline: capnography/capnometry during mechanical ventilation-2003 revision & update. *Respir Care* 2003;48(5):534-539.
62. Becker DE, Casabianca AB. Respiratory monitoring: physiological and technical considerations. *Anesth Prog* 2009;56(1):14-20.
63. Muñoz L, Morales CP, Carro JS. Capnografía, la evolución en la monitorización del paciente crítico. *Zona TES* 2013(1):16-23.
64. Bongard F, Sue D. Pulse oximetry and capnography in intensive and transitional care units. *West J Med* 1992;156(1):57-64.
65. Giner J, Casan P. Determinación de la pulsioximetría y de la capnografía en el laboratorio de función pulmonar. *Arch Bronconeumol* 2004;40(7):311-314.
66. Tavernier B, Rey D, Thevenin D, Triboulet JP, Scherpereel P. Can prolonged expiration manoeuvres improve the prediction of arterial PCO₂ from end-tidal PCO₂? *Br J Anaesth* 1997;78(5):536-540.
67. Ahmed F, Shafeeq AM, Moiz JA, Geelani MA. Comparison of effects of manual versus ventilator hyperinflation on respiratory compliance and arterial blood gases in patients undergoing mitral valve replacement. *Heart Lung* 2010;39(5):437-443.
68. Ntoumenopoulos G. Indications for manual lung hyperinflation (MHI) in the mechanically ventilated patient with chronic obstructive pulmonary disease. *Chron Respir Dis* 2005;2(4):199-207.
69. Corley A, Spooner AJ, Barnett AG, Caruana LR, Hammond NE, Fraser JF. End-expiratory lung volume recovers more slowly after closed endotracheal suctioning than after open suctioning: a randomized crossover study. *J Crit Care* 2012;27(6):742 e1-7.
70. Lachmann B. Open up the lung and keep the lung open. *Intensive Care Med* 1992;18(6):319-321.