

## Volumetric Capnography in Acute Respiratory Distress Syndrome: Is the Era of Day-to-Day Monitoring Finally Here?

Acute lung injury (ALI) and the acute respiratory distress syndrome (ARDS) affect 4.5–7.1% of the patients admitted to the intensive care unit with acute respiratory failure.<sup>1,2</sup> In our population, we found that ARDS affects 15.3 per 100,000 persons per year.<sup>3</sup> Although the reported mortality of patients with ALI and ARDS has improved, it remains high among the causes of death in the intensive care unit. Affected patients usually die secondary to sepsis and multisystem organ dysfunction. Some predictive factors associated with mortality include the triggering disease, the severity of the respiratory illness, and the presence of dysfunction of the right ventricle.<sup>4</sup> Although hypoxemia is the protean gas exchange abnormality in ALI and ARDS (it is part of the definitions),<sup>5</sup> the degree of hypoxemia does not predict outcome.<sup>6</sup> As a matter of fact, patients treated with lower tidal volume ( $V_T$ ) in the ARDS Network trial had a lower ratio of  $P_{aO_2}$  to fraction of inspired oxygen in the first 3 days of therapy but still had a lower mortality and a higher number of ventilator-free days.<sup>7</sup>

In this issue of *RESPIRATORY CARE*, Kallet et al<sup>8</sup> demonstrate that using volumetric capnography to calculate the physiologic dead space in patients with ARDS gives values that correlate with those obtained by the metabolic-cart method used in their important 2002 paper.<sup>9</sup> Kallet et al showed a statistically strong correlation between the 2 techniques in terms of the expired carbon dioxide tension and physiologic dead-space-to- $V_T$  ratio, with a fair bias and precision. However, the sample of patients was small and it will be important to independently confirm these exciting findings. These results are important because they simplify the technique of a bedside measurement that can predict survival in patients with ARDS.<sup>9</sup>

Enghoff defined the physiologic dead space in 1938 as the “wasted ventilation.”<sup>10</sup> He modified the Bohr’s equation by replacing the alveolar  $CO_2$  concentration with the  $P_{aCO_2}$ , adding to the anatomic dead space the alveolar units that are either completely nonventilated or subject to varying degrees of ventilation-perfusion mismatch. The physiologic dead space is therefore a measure of the efficiency of ventilation. In 2002, Nuckton et al identified the physiologic dead-space fraction measured within several hours of the onset of respiratory failure as the first lung-specific predictor of mortality in ARDS.<sup>9</sup> To calculate the physi-

ologic dead space they used a metabolic monitor, also known as indirect calorimetry cart, that allows the collection of the exhaled  $V_T$  in the cart’s built-in mixing chamber. An infrared sensor then measures the expired  $CO_2$  fraction in the collected gas, which in turn allows the calculation of the partial pressure of  $CO_2$  in the exhaled  $V_T$ . The technique is accurate if water vapor pressure is well controlled, the system is relatively free of leaks, and the exhaled  $V_T$  is collected for several cycles to average the breath-to-breath variation in  $CO_2$  composition. The bedside metabolic monitor has been validated against the classic method of collecting exhaled gases in a Douglas bag and measuring the  $CO_2$  partial pressure in an arterial blood gas analyzer.<sup>11</sup> The metabolic cart is less cumbersome than the Douglas bag, and it allows a minute-to-minute bedside evaluation.

---

SEE THE ORIGINAL STUDY ON PAGE 462

---

However, in our experience, the use of the metabolic cart still poses a considerable work load for the intensive care unit staff and may not be readily available, since its use for the assessment of the patient’s nutritional needs and oxygen consumption seems not to be common.

Volumetric capnography, also known as the single-breath test for  $CO_2$ , relies on a bedside instrument that displays a plot of the partial pressure of  $CO_2$  (or fractional concentration of  $CO_2$ ) in the exhaled gas as a function of the exhaled  $V_T$ . It records the partial pressure of  $CO_2$  throughout a tidal breath by acquiring a calibrated signal from an infrared light.<sup>12</sup> The light emits a wavelength near 4.28  $\mu m$ , at which the maximum absorbance of carbon dioxide occurs. The exhaled  $V_T$  is simultaneously acquired by measuring the expiratory flow that results in the familiar S-shaped  $CO_2$ - $V_T$  plot. The mean exhaled pressure of  $CO_2$  can be electronically derived and used in the calculation of the physiologic dead space. Alternatively, the arterial partial pressure of  $CO_2$  can be represented on the exhaled  $CO_2$ - $V_T$  curve as a horizontal line. The rectangular area formed by this line and the exhaled  $V_T$  represents an ideal system, free of dead space or ventilation-perfusion mismatching, where there is complete equilibration between the alveolar and arterial  $CO_2$ . The difference between this

ideal area and the area under the S-shaped curve would then be the physiologic dead space.<sup>13</sup>

Why is an elevated physiologic dead space associated with an increased mortality in patients with ARDS? Is it that it identifies a subset of patients with the “worst” degree of lung injury, as does the quasistatic respiratory compliance, another predictor of mortality? If the physiologic dead space reflects pulmonary vascular injury, does it correlate with the degree of right-ventricular dysfunction, or with levels of Von Willebrand factor, which is a marker of endothelial injury in ALI/ARDS?<sup>14,15</sup> Do patients with an increased dead-space fraction also have an impaired maximum alveolar-fluid clearance?<sup>16</sup> All of these are questions that need to be answered in the near future.

The strong statistical agreement between the metabolic cart and the volumetric capnograph implies that the latter can be accurately used to predict mortality in patients with ARDS. A simpler technique to estimate the physiologic dead space in mechanically ventilated patients should lead to important clinical and research applications. We should anticipate trials that monitor the day-to-day severity of lung injury and that potentially follow the response to therapy.

**Ziad W Ghamra MD**

**Alejandro C Arroliga MD**

Department of Pulmonary, Allergy and  
Critical Care Medicine  
The Cleveland Clinic Foundation  
Cleveland, Ohio

#### REFERENCES

1. Esteban A, Anzueto A, Frutos F, Alia I, Brochard L, Stewart TE, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA* 2002;287(3):345–355.
2. Brun-Buisson C, Minelli C, Bertolini G, Brazzi L, Pimentel J, Lewandowski K, et al. Epidemiology and outcome of acute lung injury

- in European intensive care units. *Intensive Care Med* 2004;30(1):51–61.
3. Arroliga AC, Ghamra ZW, Perez Trepichio A, Perez Trepichio P, Komara JJ Jr, Smith A, Wiedemann HP. Incidence of ARDS in an adult population of Northeast Ohio. *Chest* 2002;121(6):1972–1976.
4. Monchi M, Bellenfant F, Cariou A, Joly LM, Thebert D, Laurent I, et al. Early predictive factors of survival in the acute respiratory distress syndrome: a multivariate analysis. *Am J Respir Crit Care Med* 1998;158(4):1076–1081.
5. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, et al. The American European consensus conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149(3 Pt 1):818–824.
6. Doyle RL, Szaflarski N, Modin GW, Wiener-Kronish JP, Matthay MA. Identification of patients with acute lung injury: predictors of mortality. *Am J Respir Crit Care Med* 1995;152(6 Pt 1):1818–1824.
7. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 2000;342(18):1301–1308.
8. Kallet RH, Daniel BM, Garcia O, Matthay MA. Accuracy of physiologic dead space measurements in patients with acute respiratory distress syndrome using volumetric capnography: comparison with the metabolic monitor method. *Respir Care* 2005;50(4):462–467.
9. Nuckton TJ, Alonso JA, Kallet RH, Daniel BM, Pittet JF, Eisner MD, Matthay MA. Pulmonary dead-space fraction as a risk factor for death in the acute respiratory distress syndrome. *N Engl J Med* 2002;346(17):1281–1286.
10. Enghoff H. Volumen inefficax: Bemerkungen zur Frage des schadlichen Raumes. *Upsala Lakareforen Forh* 1938;44:191–218.
11. MacKinnon JC, Houston PL, McGuire GP. Validation of the Deltrac metabolic cart for measurement of dead-space-to-tidal-volume ratio. *Respir Care* 1997;42(8):761–764.
12. Romero PV, Lucangelo U, Lopez Aguilar J, Fernandez R, Blanch L. Physiologically based indices of volumetric capnography in patients receiving mechanical ventilation. *Eur Respir J* 1997;10(6):1309–1315.
13. Fletcher R, Johnson G, and Brew J. The concept of deadspace with special reference to the single breath test for carbon dioxide. *Br J Anesth* 1981;53:77–88.
14. Ware LB, Eisner MD, Thompson BT, Parsons PE, Matthay MA. Significance of von Willebrand factor in septic and nonseptic patients with acute lung injury. *Am J Respir Crit Care Med* 2004;170(7):766–772.
15. Robertson HT, Swenson ER. What do dead-space measurements tell us about the lung with acute respiratory distress syndrome? *Respir Care* 2004;49(9):1006–1007.
16. Ware LB, Matthay MA. Alveolar fluid clearance is impaired in the majority of patients with acute lung injury and the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001;163(6):1376–1383.

Correspondence: Alejandro C Arroliga MD, Cleveland Clinic Lerner College of Medicine, Section of Critical Care Medicine, Department of Pulmonary, Allergy and Critical Care Medicine, 9500 Euclid Avenue G-62, Cleveland OH 44195. E-mail: arrolia@ccf.org.