

AARC Clinical Practice Guideline

Removal of the Endotracheal Tube—2007 Revision & Update

RET 1.0 PROCEDURE

Elective removal of the endotracheal tube from adult, pediatric, and neonatal patients.

RET 2.0 DESCRIPTION/DEFINITION

The decision to discontinue mechanical ventilation involves weighing the risks of prolonged mechanical ventilation against the possibility of extubation failure.^{1,2} This guideline will focus on the predictors that aid the decision to extubate, the procedure referred to as extubation, and the immediate postextubation interventions that may avoid potential reintubation. This review will not address weaning from mechanical ventilation, accidental extubation, nor terminal extubation.

2.1 The risks of prolonged translaryngeal intubation include but are not limited to:

- 2.1.1** Sinusitis^{3,4}
- 2.1.2** Vocal cord injury⁵
- 2.1.3** Laryngeal injury⁶⁻⁸
- 2.1.4** Laryngeal stenosis^{6,7}
- 2.1.5** Subglottic stenosis in neonates⁹⁻¹¹ and children¹²
- 2.1.6** Tracheal injury¹³⁻¹⁶
- 2.1.7** Hemoptysis¹⁷
- 2.1.8** Aspiration^{18,19}
- 2.1.9** Pulmonary infection²⁰⁻²³
- 2.1.10** Endotracheal tube occlusion²⁴⁻²⁷
- 2.1.11** Accidental extubation necessitating emergent reintubation^{25,28}

2.2 Extubation may result in the following complications

- 2.2.1** Upper airway obstruction from laryngospasm²⁹⁻³²
- 2.2.2** Laryngeal edema³³⁻³⁷
- 2.2.3** Supraglottic obstruction³⁸
- 2.2.4** Pulmonary edema³⁹⁻⁴¹
- 2.2.5** Pulmonary aspiration syndrome^{19,42}
- 2.2.6** Impaired respiratory gas exchange

RET 3.0 ENVIRONMENT

The endotracheal tube should be removed in an environment in which the patient can be physiologically monitored and in which emergency equipment and appropriately trained health care providers with airway management skills are immediately available (see RET 10.0 and 11.0).

RET 4.0 INDICATIONS/OBJECTIVES

When the airway control afforded by the endotracheal tube is deemed to be no longer necessary for the continued care of the patient, the tube should be removed. Subjective or objective determination of improvement of the underlying condition impairing pulmonary function and/or gas exchange capacity is made prior to extubation.² To maximize the likelihood for successful extubation, the patient should be capable of maintaining a patent airway and generating adequate spontaneous ventilation. In general, this requires the patient to possess adequate: central inspiratory drive, respiratory muscle strength, cough strength to clear secretions, laryngeal function, nutritional status, and clearance of sedative and neuromuscular blocking effects.

4.1 Occasionally, acute airway obstruction of the artificial airway due to mucus or mechanical deformation mandates immediate removal of the artificial airway. Reintubation or other appropriate techniques for reestablishing the airway (ie, surgical airway management) must be used to maintain effective gas exchange.^{26,27,43}

4.2 Patients in whom an explicit declaration of the futility of further medical care is documented may have the endotracheal tube removed despite failure to meet the above indications.^{44,45}

RET 5.0 CONTRAINDICATIONS

There are no absolute contraindications to extubation; however, to maintain acceptable gas exchange

after extubation some patients may require one or more of the following: noninvasive ventilation, continuous positive airway pressure, high inspired oxygen fraction, or reintubation. Airway protective reflexes may be depressed immediately following and for some time after extubation.^{18,46} Therefore, measures to prevent aspiration should be considered.

RET 6.0 HAZARDS/COMPLICATIONS

6.1 Hypoxemia after extubation may result from but is not limited to

6.1.1 Failure to deliver adequate inspired oxygen fraction through the natural upper airway⁴⁷

6.1.2 Acute upper airway obstruction secondary to laryngospasm²⁹⁻³²

6.1.3 Development of post-obstruction pulmonary edema³⁹⁻⁴¹

6.1.4 Bronchospasm^{48,49}

6.1.5 Development of atelectasis, or lung collapse⁵⁰

6.1.6 Pulmonary aspiration^{18,19,42}

6.1.7 Hypoventilation^{51,52}

6.2 Hypercapnia after extubation may be caused by but is not limited to

6.2.1 Upper airway obstruction resulting from edema of the trachea, vocal cords, or larynx³³⁻³⁸

6.2.2 Respiratory muscle weakness^{53,54}

6.2.3 Excessive work of breathing⁵⁵⁻⁵⁹

6.2.4 Bronchospasm^{48,49}

6.3 Death may occur when medical futility is the reason for removing the endotracheal tube.

RET 7.0 LIMITATIONS OF METHODOLOGY/ VALIDATION OF RESULTS

Predicting extubation outcome is of significant clinical importance as both extubation delay and unsuccessful extubation are associated with poor patient outcomes.^{1,2} However, the literature on this topic is limited by few validated objective measures to accurately predict the extubation outcome for an individual patient.^{1,2,60-63} Failed extubation, or the need to reinsert an artificial airway following extubation, is not necessarily an indication of failed medical practice. Patients may need reintubation immediately or after some interval due to inappropriately timed extubation, progression of underlying disease, or development of a new disorder.

Therefore, a trial of extubation may be used in some marginal patients with the expectation that the need for reintubation is likely. Extubation failure rates reported in the literature range between 1.8%-18.6% for adults,^{1,36,63-65} 2.7%-22% for children,⁶⁶⁻⁷⁰ and may be as high as 40%-60% for low birth-weight infants.⁷¹⁻⁷⁵ Clinical practice standards for endotracheal tube removal include attentive postextubation monitoring, prompt identification of respiratory distress, maintenance of a patent airway and, if clinically indicated, attempts to successfully establish an artificial airway by reintubation or surgical technique. The failure and complication rates of extubation can be used as quality monitors.

RET 8.0 ASSESSMENT OF EXTUBATION READINESS

The endotracheal tube should be removed as soon as the patient no longer requires an artificial airway. Patients should demonstrate some evidence for the reversal of the underlying cause of respiratory failure and should be capable of maintaining adequate spontaneous ventilation and gas exchange. The determination of extubation readiness may be individualized using the following guidelines.

8.1 Patients with an artificial airway to facilitate treatment of respiratory failure should be considered for extubation when they have met established extubation readiness criteria.^{64,76} Examples of these criteria include but are not limited to

8.1.1 The capacity to maintain adequate arterial partial pressure of oxygen (P_{aO_2}/F_{IO_2} ratio > 150-200) on inspired oxygen fractions provided with simple oxygen devices ($F_{IO_2} \leq 0.4$ to 0.5 and with low levels of positive airway pressure (PEEP) ≤ 5 to 8 cm H₂O²

8.1.2 The capacity to maintain appropriate pH (pH ≥ 7.25)² and arterial partial pressure of carbon dioxide during spontaneous ventilation^{74,75}

8.1.3 Successful completion of 30-120 minute spontaneous breathing trial (SBT) performed with a low level of CPAP (eg 5 cm H₂O) or low level of pressure support (eg 5-7 cm H₂O) demonstrating adequate respiratory pattern and gas exchange, hemodynamic stability, and subjective

comfort⁷⁷⁻⁸¹

8.1.4 In adults, respiratory rate < 35 breaths per minute during spontaneous breathing;⁵⁴ in infants and children, the acceptable respiratory rate decreases inversely with age and can be measured with good repeatability with a stethoscope⁸²

8.1.5 Adequate respiratory muscle strength⁸³⁻⁸⁵

8.1.6 Maximum negative inspiratory pressure > -30 cm H₂O⁸⁶⁻⁸⁹ although current clinical practice may accept a maximum negative inspiratory pressure > -20 cm H₂O^{90,91}

8.1.7 Vital capacity > 10 mL/kg ideal body weight⁹² or in neonates > 150 mL/m²⁸⁹

8.1.8 Pressure measured across the diaphragm during spontaneous ventilation < 15% of maximum^{93,94}

8.1.9 In adults, spontaneous exhaled minute ventilation < 10 L/min⁸⁶

8.1.10 In adults, a rapid shallow breathing index (RSBI, respiratory rate-to-tidal-volume ratio) of ≤105 breaths/min (positive predictive value (PPV) of 0.78)⁹⁰; in infants and children, variables standardized by age or weight proved more useful. Modified CROP index (compliance, resistance, oxygenation, and ventilating pressure) above a threshold of ≥ 0.1-0.15 mL · mmHg/breaths/min/kg (sensitivity of 83% and specificity of 53%) may be a superior screening tool than a modified RSBI ≤ 8-11 breaths/min/mL/kg (sensitivity of 74% and specificity of 74%)^{67,68,95}

8.1.11 Thoracic compliance > 25 mL/cm H₂O⁹⁶

8.1.12 Work of breathing < 0.8 J/L⁹⁷⁻¹⁰²

8.1.13 Oxygen cost of breathing < 15% total especially for those patients with chronic respiratory insufficiency requiring long-term mechanical ventilation (sensitivity, 100%; specificity, 87%)¹⁰²⁻¹⁰⁴

8.1.14 Dead-space-to-tidal-volume ratio (V_D/V_T) < 0.6; in children, V_D/V_T ≤ 0.5 equates to 96% successful extubation, 0.51-0.64 equates to 60% successful extu-

bation, (0.65 equates to 20% successful extubation)^{105,106}

8.1.15 Airway occlusion pressure at 0.1 seconds ($P_{0.1}$) < 6 cm H₂O and when normalized for maximal inspiratory pressure (MIP), as indicated by [$P_{0.1}/MIP$], accurately predict successful extubation 88% and 98% of the time, respectively.¹⁰⁷⁻¹⁰⁹ (This measurement is primarily a research tool.)

8.1.16 Maximum voluntary ventilation > twice resting minute ventilation⁸⁶

8.1.17 In preterm infants, minute ventilation testing vs. standard clinical evaluation resulted in shorter time to extubation¹¹⁰

8.1.19 Peak expiratory flow (PEF) ≥ 60 L/min after 3 cough attempts measured with an in-line spirometer^{111,112}

8.1.20 Time to recovery of minute ventilation to pre-spontaneous breathing trial baseline levels¹¹³

8.1.21 Sustained maximal inspiratory pressures (SMIP) > 57.5 pressure time units (sensitivity and specificity of 1.0) predicted extubation outcome¹¹⁴

8.1.22 In neonates, total respiratory compliance (C_{RS} , derived from $V_T/PIP-PEEP$) ≤ 0.9 mL/cm H₂O was associated with extubation failure, whereas a value ≥ 1.3 mL/cm H₂O was associated with extubation success¹¹⁵

8.1.23 Preterm infants extubated directly from low rate ventilation without a trial of endotracheal tube continuous positive airway pressure (CPAP) demonstrated a trend to increased chance of successful extubation (RR = 0.45, CI₉₅ (0.19, 1.07), NNT 10)¹¹⁶

8.1.24 Integrated indices of measured vital capacity (VC, threshold value = 635 mL), respiratory frequency to tidal volume ratio (f/V_T , threshold value = 88 breaths/min/L) and maximal expiratory pressure (MEP, threshold value = 28 cm H₂O)^{80,117}

8.2 In addition to treatment of respiratory failure, artificial airways are sometimes placed for airway protection. Resolution of the need for airway protection may be assessed by but is not

limited to

8.2.1 Appropriate level of consciousness¹¹⁸⁻¹²⁰

8.2.2 Adequate airway protective reflexes^{119,120}

8.2.2.1 Reduced cough strength (grade 0 to 2) measured by the white card test and increased secretion burden predicted unsuccessful extubation (risk ratio, RR= 4.0; 95% confidence interval (CI₉₅) 1.8 to 8.9)¹²¹

8.2.3 Easily managed secretions^{1,63,119,120}

8.3 In addition to resolution of the processes requiring the insertion of an artificial airway, issues that should be considered in all patients prior to extubation are

8.3.1 No immediate need for reintubation anticipated

8.3.2 Known risk factors for extubation failure

8.3.2.1 Patient features of high risk for extubation failure include: admission to medical ICU, age > 70 or < 24 months, higher severity of illness upon weaning, Hgb <10 mg/dL, use of continuous I.V. sedation, longer duration of mechanical ventilation, presence of a syndromic or chronic medical condition, known medical or surgical airway condition^{63,70}, frequent pulmonary toilet,¹²¹ and loss of airway protective reflexes^{119,120}

8.3.2.2 Risk factors for a known history of a difficult airway: syndromic or congenital conditions associated with cervical instability (ie, Klippel-Feil or Trisomy 21); limited physical access to the airway (ie, halo-vest or anatomic hindrances); multiple failed direct laryngoscopy attempts by an experienced laryngoscopist, or a failed laryngoscopy attempt followed by tracheal intubation using fiberoptic bronchoscopy or nasal lightwand, or requiring placement of a laryngeal mask airway¹²²⁻¹²⁹

8.3.2.3 In the pediatric cardiothoracic surgery population, presence of one or more of these variables increases the likelihood of failed extubation: age < 6

months, history of prematurity, congestive heart failure, and pulmonary hypertension¹³⁰

8.3.2.4 For pediatric patients, validated bedside measures of respiratory function identifying low (< 10%) and high risk (> 25%) threshold values of extubation failure which may be useful in generating discussion, but do not apply to individual risk¹³¹

8.3.3 Presence of upper airway obstruction or laryngeal edema as detected by diminished gas leak around the endotracheal tube with positive pressure breaths^{37,132-139}

8.3.3.1 Percent cuff leak or the difference between expiratory tidal volume measured with the cuff inflated and then deflated in a volume-controlled mode of $\geq 15.5\%$ (sensitivity 75%, specificity 72%).¹³²⁻¹³⁸ Yet this test was found not to be predictive in a study of cardiothoracic surgery patients.¹³⁹

8.3.3.2 Air leak may be an age-dependent predictor of postextubation stridor in children. An air leak > 20 cm H₂O was predictive of postextubation stridor in children ≥ 7 years of age (sensitivity of 83%, specificity of 80%), but was not predictive < 7 years of age³⁷

8.3.3.3 Air leak test has been predictive of postextubation stridor or extubation failure for children with upper airway pathology: trauma patients,¹⁴⁰ croup,¹⁴¹ after tracheal surgery.¹⁴²

8.3.4 Evidence of stable, adequate hemodynamic function^{2,143-146}

8.3.5 Evidence of stable nonrespiratory functions¹⁴⁷⁻¹⁵⁰

8.3.6 Electrolyte values within normal range¹⁵¹⁻¹⁵³

8.3.7 Evidence of malnutrition decreasing respiratory muscle function and ventilatory drive¹⁵⁴⁻¹⁵⁸

8.3.8 Anesthesia literature indicates the patient must have no intake of food or liquid by mouth for a period of time prior to airway manipulation. The continuation of transpyloric feedings during an extubation procedure remains controversial^{159,160}

8.3.9 Prophylactic medication prior to extubation to avoid or reduce the severity of postextubation complications such as

8.3.9.1 Consider use of lidocaine to prevent cough and/or laryngospasm in patients at risk^{161,162}

8.3.9.2 Prophylactic administration of steroids may be helpful to prevent re-intubation rates in high-risk neonates, but not in children (relative risk, RR = 0.49, CI₉₅ 0.01, 19.65) or adults (RR = 0.95, CI₉₅ 0.52, 1.72)^{61,163}

8.3.9.3 Prophylactic administration of steroids may help reduce the incidence of postextubation stridor in children (RR = 0.53, CI₉₅ 0.28, 0.97), but not in neonates or adults (RR = 0.86, CI₉₅ 0.57, 1.30)^{61,163}

8.3.9.4 Prophylactic administration of steroids for patients with laryngotracheobronchitis (croup) correlates with reduced rates of reintubation^{164,165}

8.3.9.5 Caffeine citrate reduced the risk of apnea for infants, but did not reduce the risk of extubation failure¹⁶⁶

8.3.9.6 Methylxanthine treatment stimulates breathing and reduces the rate of apnea (RR = 0.47, CI₉₅ (0.32, 0.70). [Number needed to treat (NNT) 3.7, CI₉₅ (2.7, 6.7)] for neonates with poor respiratory drive, especially extremely low birthweight infants.¹⁶⁷

RET 9.0 ASSESSMENT OF OUTCOME

Removal of the endotracheal tube should be followed by adequate spontaneous ventilation through the natural airway, adequate oxygenation, and no need for re-intubation.

9.1 Clinical outcome may be assessed by physical examination, auscultation, invasive and noninvasive measurements of gas exchange, and chest radiography.

9.2 Quality of the procedure can be systematically assessed by monitoring extubation complications and the need for reintubation.

9.3 The success of removal of the endotracheal tube can be monitored by examining the frequency of reintubation and frequency of complications.

9.4 When a patient experiences an unplanned

self-extubation and does not require reintubation, this suggests that planned extubation should have been considered earlier.¹⁶⁸⁻¹⁷³

9.5 Some patients may require postextubation support or intervention to maintain adequate gas exchange independent of controlled mechanical ventilation.

9.5.1 Noninvasive Respiratory Support

9.5.1.1 Infants extubated to nasal intermittent positive pressure ventilation (NIPPV) were less likely to fail extubation than those infants extubated to nasal continuous positive airway pressure ventilation (NCPAP)[RR = 0.21, CI₉₅ (0.10, 0.45); NNT 3, CI₉₅ (2, 5)]⁵²

9.5.1.2 In neonates and premature infants, binasal prong CPAP is more effective at preventing re-intubation than single nasal or nasopharyngeal prongs [RR = 0.59, CI₉₅ (0.41, 0.85); NNT 5, CI₉₅ (3, 14)]¹⁷⁴

9.5.1.3 In adults, routine use of postextubation noninvasive positive pressure ventilation such as BIPAP is not supported¹⁷⁵⁻¹⁷⁷

9.5.1.4 In patients with chronic obstructive pulmonary disease (COPD), continuous positive airway pressure (CPAP) of 5 cm H₂O and pressure support ventilation (PSV) of 15 cm H₂O have improved pulmonary gas exchange, decreased intrapulmonary shunt fraction, and reduced patient work of breathing¹⁷⁸

9.5.2 Postextubation Medical Therapy

9.5.2.1 Aerosolized levo-epinephrine is as effective as aerosolized racemic epinephrine in the treatment of postextubation laryngeal edema in children³⁵

9.5.2.2 No randomized studies in neonates have been performed to evaluate the role of nebulized racemic epinephrine for postextubation stridor¹⁷⁹

9.5.2.3 Heliox may alleviate the symptoms of partial airway obstruction and resultant stridor, improve patient comfort, decrease work of breathing, and prevent reintubation^{57,58,140}

9.5.3 Diagnostic Therapy

9.5.3.1 For patients with postextubation complications such as stridor or obstruction, fiberoptic bronchoscopy may provide direct airway inspection and therapeutic interventions (secretion clearance, instillation of drugs, removal of aspirated foreign objects).^{180,181}

RET 10.0 RESOURCES

Preparation for extubation includes assuring emergency reintubation equipment and personnel are readily available. The following equipment/supplies must be maintained in close proximity to the patient, in sufficient quantities, and in working condition.

10.1 Equipment

- 10.1.1** Oxygen source
- 10.1.2** Devices to deliver oxygen-enriched gas mixtures
- 10.1.3** High-volume suction source and catheters
- 10.1.4** Pharyngeal and tracheal suction catheters
- 10.1.5** Self-inflating or nonself-inflating manual ventilation system
- 10.1.6** Appropriately sized face masks
- 10.1.7** Oral and nasopharyngeal airways
- 10.1.8** Endotracheal tubes of various sizes, cuffed and uncuffed
- 10.1.9** Translaryngeal intubation equipment (laryngoscope blades, handles, extra batteries, stylettes, surgical lubricant, syringes to inflate cuff)
- 10.1.10** Airway exchange catheter of various sizes
- 10.1.11** Laryngeal mask airway (LMA) of various sizes
- 10.1.12** Equipment for establishing an emergency surgical airway (scalpel, lidocaine with epinephrine, appropriately sized endotracheal or tracheostomy tubes)
- 10.1.13** Nasogastric tubes of various sizes
- 10.1.14** Pulse oximeter
- 10.1.15** Two-channel cardiac monitor
- 10.1.16** Supplies for arterial puncture and blood gas analysis
- 10.1.17** Medication for sedation, analgesia, neuromuscular blockade and prevention of raised intracranial pressure as indi-

cated by the individual situation

10.1.18 Carbon dioxide detection devices (qualitative and/or quantitative devices)

10.2 Personnel

10.2.1 Credentialed and/or licensed health care personnel with documented knowledge and demonstrated skills specific to patient assessment and airway management should determine the appropriateness of extubation, be available to assess success, and begin appropriate interventions should immediate complications occur. Personnel skilled in endotracheal intubation and the insertion of invasive airways should be immediately available whenever extubation is performed.

10.2.2 Credentialed and/or licensed health-care personnel with documented knowledge and demonstrated skill in providing oxygen administration devices and suctioning the airway may provide support during the extubation procedure.

10.2.3 In the event of acute obstruction of the artificial airway, anyone with airway maintenance skills may remove the endotracheal tube to save the patient's life¹⁶⁰

RET 11.0 MONITORING

Monitoring in the postextubation period includes ensuring the equipment, personnel, and medications are readily available in the event of emergent, postextubation phenomena.¹⁶⁰

11.1 Appropriately trained personnel need to be readily available to detect cardiopulmonary impairment

11.2 Frequent respiratory evaluation should include: vital signs, assessment of neurologic status, patency of airway, auscultatory findings, work of breathing, and hemodynamic status

11.3 Equipment

11.2.1 Pulse oximeter

11.2.2 Two-channel cardiac monitor

11.2.3 Sphygmomanometer and stethoscope

11.2.4 Capnograph

RET 12.0 FREQUENCY

No consensus exists on the appropriate timing of or requirement for tracheostomy placement in the mechanically ventilated patient. Any recommendation

will have to consider patient population, etiology of respiratory insult, expected or known duration of mechanical ventilation, balance of risks and perceived benefits of continued mechanical ventilation via tracheostomy as opposed to a translaryngeally-placed endotracheal tube. Past recommendations have been based upon expert consensus.^{2,182,183}

12.1 Limited data exist on the rate of successful extubation after a previous failed extubation.

12.1.1 Many clinical studies include the first extubation attempt only

12.1.2 In a pediatric descriptive study, 174/2794 subjects failed extubation [extubation failure rate 6.2%, CI₉₅ 5.3,7.1] after the first attempt; 27% (65/174) failed a second extubation attempt; of those patients, 22 extubated successfully after the third extubation attempt.⁷⁰

RET 13.0 STANDARD PRECAUTIONS

Caregivers should exercise Standard Precautions for all patients, follow the Centers for Disease Control and Prevention (CDC) recommendations for control of exposure to tuberculosis and droplet nuclei, and institute appropriate precautions empirically for airborne, droplet, and contact agents pending confirmation of diagnosis in patients suspected of having serious infections.¹⁸⁴⁻¹⁸⁹

Revised by Angela T Wratney MD and Ira M Cheifetz MD FAARC, Duke University Medical Center, Durham, North Carolina, and approved by the 2006 CPG Steering Committee

Original publication: *Respir Care* 1999;44(1):85-90.

REFERENCES

1. Epstein SK. Decision to extubate. *Intensive Care Med* 2002;28(5):535-546.
2. MacIntyre NR, Cook DJ, Ely EW Jr, Epstein SK, Fink JB, Heffner JE, et al. Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College of Critical Care Medicine. *Chest* 2001;120(6 Suppl):375S-395S.
3. Holzapfel L, Chevret S, Madinier G, Ohen F, Demingon G, Coupary A, Chaudet M. Influence of long-term, oro- or nasotracheal intubation on nosocomial maxillary sinusitis and pneumonia: results of a prospective, randomized, clinical trial. *Crit Care Med* 1993;21(8):1132-1138.
4. Guerin JM, Lustman C, Meyer P, Barbotin-Larrieau F. Nosocomial sinusitis in pediatric intensive care patients (comment). *Crit Care Med* 1990;18(8):902.
5. Kastanos N, Estopa Miro R, Marin Perez A, Xaubet Mir A, Agusti-Vidal A. Laryngotracheal injury due to endotracheal intubation: incidence, evolution, and predisposing factors: a prospective long-term study. *Crit Care Med* 1983;11(5):362-367.
6. Colice GL, Stukel TA, Dain B. Laryngeal complications of prolonged intubation. *Chest* 1989;96(4):877-884.
7. Santos PM, Afrassiabi A, Weymuller EA Jr. Risk factors associated with prolonged intubation and laryngeal injury. *Otolaryngol Head Neck Surg* 1994;111(4):453-459.
8. Benjamin B. Prolonged intubation injuries of the larynx: endoscopic diagnosis, classification, and treatment. *Ann Otol Rhinol Laryngol* 1993;160:1-15.
9. Sherman JM, Lowitt S, Stephenson C, Ironson G. Factors influencing acquired subglottic stenosis in infants. *J Pediatrics* 1986;109(2):322-327.
10. Marcovich M, Pollauf F, Burian K. Subglottic stenosis in newborns after mechanical ventilation. *Prog Pediatr Surg* 1987;21:8-19.
11. Parkin JL, Stevens MH, Jung AL. Acquired and congenital subglottic stenosis in the infant. *Ann Otol Rhinol Laryngol* 1976; 85(5 Pt1): 573-581.
12. Wiel E, Vilette B, Darras JA, Scherpereel P, Leclerc F. Laryngotracheal stenosis in children after intubation. Report of five cases. *Paediatr Anaesthes* 1997;7(5):415-419.
13. Stauffer JL, Olson DE, Petty TL. Complications and consequences of endotracheal intubation and tracheotomy: a prospective study of 150 critically ill adult patients. *Am J Med* 1981;70(1):65-76.
14. Hoeve LJ, Eskici O, Verwoerd CD. Therapeutic reintubation for post-intubation laryngotracheal injury in preterm infants. *Int J Pediatr Otorhinolaryngol* 1995;31(1):7-13.
15. Stauffer JL, Silvestri RE. Complications of endotracheal intubation, tracheostomy, and artificial airways. *Respir Care* 1982;27(4):417-434.
16. Spear RM, Sauder RA, Nichols DG. Endotracheal tube rupture, accidental extubation, and tracheal avulsion: three airway catastrophes associated with significant decrease in leak pressure. *Crit Care Med* 1989;17(4):701-703.
17. Keceligil HT, Erk MK, Kolbakir F, Yildirim A, Yilman M, Unal R. Tracheoinnominate artery fistula following tracheostomy. *Cardiovasc Surg* 1995;3(5):509-510.
18. Leder SB, Cohn SM, Moller BA. Fiberoptic endoscopic documentation of the high incidence of aspiration following extubation in critically ill trauma patients. *Dysphagia* 1998;13(4):208-212.
19. Goitein KJ, Rein AJ, Gornstein A. Incidence of aspiration in endotracheally intubated infants and children. *Crit Care Med* 1984;12(1):19-21.
20. Garibaldi RA, Britt MR, Coleman ML, Reading JC,

- Pace NL. Risk factors for postoperative pneumonia. *Am J Med* 1981;70(3):677-680.
21. Kerver AJ, Rommes JH, Mevissen-Verhage EA, Hulstaert PF, Vos A, Verhoef J, Wittebol P. Prevention of colonization and infection in critically ill patients: a prospective randomized study. *Crit Care Med* 1988;16(11):1087-1093.
 22. Torres A, Aznar R, Gatell JM, Jiminez P, Gonzalez J, Ferrer A, et al. Incidence, risk, and prognosis factors of nosocomial pneumonia in mechanically ventilated patients. *Am Rev Respir Dis* 1990;142(3):523-528.
 23. Harris H, Wirtschafter D, Cassady G. Endotracheal intubation and its relationship to bacterial colonization and systemic infection of newborn infants. *Pediatrics* 1976;58(6):816-823.
 24. Rivera R, Tibbals J. Complications of endotracheal intubation and mechanical ventilation in infants and children. *Crit Care Med* 1992;20(2):193-199.
 25. Khan FH, Khan FA, Irshad R, Kamal RS. Complications of endotracheal intubation in mechanically ventilated patients in a general intensive care unit. *J Pak Med Assoc* 1996;46(9):195-198.
 26. Cohen IL, Weinberg PF, Fein IA, Rowinski GS. Endotracheal tube occlusion associated with the use of heat and moisture exchangers in the intensive care unit. *Crit Care Med* 1988;16(3):277-279.
 27. Skoulas IG, Kountakis SE. Endotracheal tube obstruction: a rare complication in laser ablation of recurrent laryngeal papillomas. *Ear Nose Throat J* 2003;82(7):504-506, 512.
 28. Black AE, Hatch DJ, Nauth-Misir N. Complications of naotracheral intubation in neonates, infants and children: a review of 4 years' experience in a children's hospital. *Br J Anaesth* 1990;65(4):461-467.
 29. Backus WW, Ward RR, Vitkun SA, Fitzgerald D, Askanazi J. Postextubation laryngeal spasm in an unanesthetized patient with Parkinson's disease. *J Clin Anesth* 1991;3(4):314-316.
 30. Guffin TN, Har-el G, Sanders A, Lucente FE, Nash M. Acute postobstructive pulmonary edema. *Otolaryngol Head Neck Surg* 1995;112(2):235-237.
 31. Wilson GW, Bircher NG. Acute pulmonary edema developing after laryngospasm: report of a case. *J Oral Maxillofac Surg* 1995;53(2):211-214.
 32. Young A, Skinner TA. Laryngospasm following extubation in children (letter). *Anaesthesia* 1995;50(9):827.
 33. Hartley M, Vaughan RS. Problems associated with tracheal extubation. *Br J Anaesth* 1993;71(4):561-568.
 34. Darmon JY, Rauss A, Dreyfuss D, Bleichner G, Elkharrat D, Schlemmer B, et al. Evaluation of risk factors for laryngeal edema after tracheal extubation in adults and its prevention by dexamethasone: a placebo-controlled, double-blind, multicenter study. *Anesthesiology* 1992;77(2):245-251.
 35. Nutman J, Brooks LJ, Deakins KM, Baldesare KK, Witte MK, Reed MD. Racemic versus l-epinephrine aerosol in the treatment of postextubation laryngeal edema: results from a prospective, randomized, double-blind study. *Crit Care Med* 1994;22(10):1591-1594.
 36. Kemper KJ, Benson MS, Bishop MJ. Predictors of postextubation stridor in pediatric trauma patients. *Crit Care Med* 1991;19(3):352-355.
 37. Mhanna MJ, Zamel YB, Tichy CM, Super DM. The "air leak" test around the endotracheal tube, as a predictor of postextubation stridor, is age dependent in children. *Crit Care Med* 2002;30(12):2639-2643.
 38. Vauthy PA, Reddy R. Acute upper airway obstruction in infants and children: evaluation by the fiberoptic bronchoscope. *Ann Otol Rhinol Laryngol* 1980;89(5 Pt 1):417-418.
 39. Willms D, Shure D. Pulmonary edema due to upper airway obstruction in adults. *Chest* 1988;94(5):1090-1092.
 40. Kamal RS, Agha S. Acute pulmonary oedema: a complication of upper airway obstruction. *Anaesthesia* 1984;39(5):464-467.
 41. Guinard JP. Laryngospasm-induced pulmonary edema. *Int J Pediatr Otorhinolaryngol* 1990;20(2):163-168.
 42. Arandia HY, Grogono AW. Comparison of the incidence of combined "risk factors" for gastric acid aspiration: influence of two anesthetic techniques. *Anesth Analg* 1980;59(11):862-864.
 43. Kapadia FN, Bajan KB, Singh S, Mathew B, Nath A, Wadkar S. Changing patterns of airway accidents in intubated ICU patients. *Intensive Care Med* 2001;27(1):296-300.
 44. Task Force on Ethics of the Society of Critical Care Medicine. Consensus report on the ethics of foregoing life-sustaining treatments in the critically ill. *Crit Care Med* 1990;18(12):1435-1439.
 45. Truog RD, Cist AF, Brackett SE, Burns JP, Curley MA, Danis M, et al. Recommendations for end-of-life care in the intensive care unit: The Ethics Committee of the Society of Critical Care Medicine. *Crit Care Med* 2001;29(12):2332-2348.
 46. Ajemian MS, Nirmul GB, Anderson MT, Zirlen DM, Kwasnik EM. Routine fiberoptic endoscopic evaluation of swallowing following prolonged intubation: implications for management. *Arch Surg* 2001;136(4):434-437.
 47. Guglielminotti J, Constant I, Murat I. Evaluation of routine tracheal extubation in children: inflating or suctioning technique? *Br J Anaesth* 1998;81(5):692-695.
 48. Tait AR, Malviya S, Voepel-Lewis T, Munro HM, Seiwert M, Pandit UA. Risk factors for perioperative adverse respiratory events in children with upper respiratory tract infections. *Anesthesiology* 2001;95(2):299-306.
 49. Wong DH, Weber EC, Schell MJ, Wong AB, Anderson CT, Barker SJ. Factors associated with postoperative pulmonary complications in patients with severe chronic obstructive pulmonary disease. *Anesth Analg* 1995;80(2):276-284.
 50. Flenady VJ, Gray PH. Chest physiotherapy for preventing morbidity in babies being extubated from mechanical ventilation. *Cochrane Database Syst Rev* 2002;(2):CD000283.

51. Lopata M, Onal E. Mass loading, sleep apnea, and the pathogenesis of obesity hypoventilation. *Am Rev Respir Dis* 1982;126(4):640-645.
52. Davis PG, Lemyre B, de Paoli AG. Nasal intermittent positive pressure ventilation (NIPPV) versus nasal continuous positive airway pressure (NCPAP) for preterm neonates after extubation. *Cochrane Database Syst Rev* 2001;(3):CD003212.
53. Mier A, Laroche C, Agnew JE, Vora H, Clarke SW, Green M, Pavia D. Tracheobronchial clearance in patients with bilateral diaphragmatic weakness. *Am Rev Respir Dis* 1990;142(3):545-548.
54. Cohen CA, Zigelbaum G, Gross D, Roussos C, Macklem PT. Clinical manifestations of inspiratory muscle fatigue. *Am J Med* 1982;73(3):308-316.
55. Sundaram RK, Nikolic G. Successful treatment of post-extubation stridor by continuous positive airway pressure. *Anesthes Intensive Care* 1996;24(3):392-393.
56. Hertzog JH, Siegel LB, Hauser GJ, Dalton HJ. Noninvasive positive-pressure ventilation facilitates tracheal extubation after laryngotracheal reconstruction in children. *Chest* 1999;116(1):260-263.
57. Kemper KJ, Izenberg S, Marvin JA, Heimbach DM. Treatment of postextubation stridor in a pediatric patient with burns: the role of heliox. *J Burn Care Rehabil* 1990;11(4):337-339.
58. Kemper KJ, Ritz RH, Benson MS, Bishop MS. Helium-oxygen mixture in the treatment of postextubation stridor in pediatric trauma patients. *Crit Care Med* 1991;19(3):356-359.
59. Jaber S, Carlucci A, Boussarsar M, Fodil R, Pigeot J, Maggiore S, et al. Helium-oxygen in the postextubation period decreases inspiratory effort. *Am J Respir Crit Care Med* 2001;164(4):633-637.
60. Epstein SK. Endotracheal extubation. *Respir Care Clin N Am* 2000;6(2):321-360.
61. Davis PG, Henderson-Smart DJ. Intravenous dexamethasone for extubation of newborn infants. *Cochrane Database Syst Rev* 2001;(4):CD000308.
62. Sinha SK, Donn SM. Weaning newborns from mechanical ventilation. *Semin Neonatol* 2002;7(5):421-428.
63. Epstein SK. Predicting extubation failure: is it in (on) the cards? (editorial) *Chest* 2001;120(4):1061-1063.
64. DeHaven CB Jr, Hurst JM, Btanson RD. Evaluation of two different extubation criteria: attributes contributing to success. *Crit Care Med* 1986;14(2):92-94.
65. Ely EW, Baker AM, Dunagan DP, Burke HL, Smith AC, Kelly PT, et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med* 1996;335(25):1864-1869.
66. el-Khatib MF, Baumeister B, Smith PG, Chatburn RL, Blumer JL. Inspiratory pressure/maximal inspiratory pressure: does it predict successful extubation in critically ill infants and children? *Intensive Care Med* 1996;22(3):264-268.
67. Farias JA, Alia I, Esteban A, Golubicki AN, Olazarri FA. Weaning from mechanical ventilation in pediatric intensive care patients. *Intensive Care Med* 1998;24(10):1070-1075.
68. Baumeister BL, el-Khatib M, Smith PG, Blumer JL. Evaluation of predictors of weaning from mechanical ventilation in pediatric patients. *Pediatr Pulmonol* 1997;24(5):344-352.
69. Khan N, Brown A, Venkataraman ST. Predictors of extubation success and failure in mechanically ventilated infants and children. *Crit Care Med* 1996; 24(9):1568-1579.
70. Kurachek SC, Newth CJ, Quasney MW, Rice T, Sachdeva RC, Patel NR, et al. Extubation failure in pediatric intensive care: a multiple-center study of risk factors and outcomes. *Critical Care Medicine* 2003;31(11):2657-2664.
71. Allatar, MH, Bhola M, Weiss M, Dallal O, Muraskas JK. The micropreemies: how early could they be extubated? *Pediatric Research* 2001;49:360A.
72. Sillos EM, Veber M, Schulman M, Krauss AN, Auld PA. Characteristics associated with successful weaning in ventilator-dependent preterm infants. *Am J Perinatol* 1992; 9(5-6):374-377.
73. Veness-Meehan K, Richter S, Davis JM. Pulmonary function testing prior to extubation in infants with respiratory distress syndrome. *Pediatr Pulmonol* 1990;9(1):2-6.
74. Berman LS, Fox WW, Raphaely RC, Downes JJ Jr. Optimum levels of CPAP for tracheal extubation of newborn infants. *J Pediatr* 1976;89(1):109-112.
75. Kim EH, Boutwell WC. Successful direct extubation of very low birthweight infants from low intermittent mandatory ventilation rate. *Pediatrics* 1987; 80(3):409-414. *Erratum in: Pediatrics* 1987;80(6):948.
76. Leitch EA, Moran JL, Greally B. Weaning and extubation in the intensive care unit: clinical or index-driven approach? *Intensive Care Med* 1996;22(8):752-759.
77. Farias JA, Retta A, Alia I, Olazarri F, Esteban A, Golubicki A, et al. A comparison of two methods to perform a breathing trial before extubation in pediatric intensive care patients. *Intensive Care Med* 2001;27(10):1649-1654.
78. Esteban A, Alia I, Gordo F, Fernandez R, Solsona JF, Vallverdu I, et al. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. The Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1997;156(2 Pt 1):459-465. *Erratum in: Am J Respir Crit Care Med* 1997;156(6):2028.
79. Esteban A, Alia I, Tobin MJ, Gil A, Gordo F, Vallverdu I, et al. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. The Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1999;159(2):512-518.
80. Vallverdu I, Calaf N, Subirana M, Net A, Benito S, Mancebo J. Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation.

- Am J Respir Crit Care Med 1998;158(6):1855-1862.
81. Ely EW, Baker AM, Evans GW, Haponik EF. The prognostic significance of passing a daily screen of weaning parameters. *Intensive Care Med* 1999;25(6):581-587.
 82. Rusconi F, Castagneto M, Gagliardi L, Leo G, Pellegratta A, Porta N, et al. Reference values for respiratory rate in the first 3 years of life. *Pediatrics* 1994;94(3):350-355.
 83. Bellemare F, Grassino A. Effect of pressure and timing of contraction on human diaphragm fatigue. *J Appl Physiol* 1982;53(5):1190-1195.
 84. Jubran A, Tobin MJ. Passive mechanics of lung and chest wall in patients who failed or succeeded in trials of weaning. *Am J Respir Crit Care Med* 1997;155(3):916-921.
 85. Roussos C, Macklem PT. The respiratory muscles. *N Engl J Med* 1982;307(13):786-797.
 86. Sahn SA, Lakshminarayan S. Bedside criteria for discontinuation of mechanical ventilation. *Chest* 1973;63(6):1002-1005.
 87. Hess D. Measurement of maximal inspiratory pressure: a call for standardization (editorial). *Respir Care* 1989;34(10):857-559.
 88. Marini JJ, Smith TC, Lamb V. Estimation of inspiratory muscle strength in mechanically ventilated patients: the measurement of maximal inspiratory pressure. *J Crit Care* 1986;1(1):32-38.
 89. Belani KG, Gilmour IJ, McComb C, Williams A, Thompson TR. Preextubation ventilatory measurements in newborns and infants. *Anesth Analg* 1980;59(7):467-472.
 90. Yang KL, Tobin MJ. A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991;324(21):1445-1450.
 91. Martin L, Bratton S, Walker L. Principles and practice of respiratory support and mechanical ventilation. In: Rogers MC, editor. *Textbook of pediatric intensive care*, 3rd ed. Baltimore: Williams & Williams; 1996.
 92. Bendixen HH, Egbert LD, Hedly-White J, et al. Management of patients undergoing prolonged artificial ventilation. *Respir Care* 1965;10:149-153.
 93. Aubier M, Murciano D, Lecoquic Y, Viires N, Pariente R. Bilateral phrenic stimulation: a simple technique to assess diaphragmatic fatigue in humans. *J Appl Physiol* 1985;58(1):58-64.
 94. Tobin MJ, Laghi F, Jubran A. Respiratory muscle dysfunction in mechanically-ventilated patients. *Mol Cell Biochem* 1998;179(1-2):87-98.
 95. Thiagarajan RR, Bratton SL, Martin LD, Brogan TV, Taylor D. Predictors of successful extubation in children. *Am J Respir Crit Care Med* 1999;160(5 Pt 1):1562-1566.
 96. Peters RM, Hilberman M, Hogan JS, Crawford DA. Objective indications for respiratory therapy in post-trauma and postoperative patients. *Am J Surg* 1972;124(2):262-269.
 97. Lee KH, Hui LP, Chan TB, Tan WC, Lim TK. Rapid shallow breathing (frequency-tidal volume ratio) did not predict extubation outcome. *Chest* 1994;105(2):540-543.
 98. Krieger BP, Isber J, Breitenbucher A, Throop G, Ershowsky P. Serial measurements of the rapid-shallow-breathing index as a predictor of weaning outcome in elderly medical patients. *Chest* 1997;112(4):1029-1034.
 99. Lewis WD, Chwals W, Benotti PN, Lakshman K, O'Donnell C, Blackburn GL, Bistrrian BR. Bedside assessment of the work of breathing. *Crit Care Med* 1988;16(2):117-122.
 100. Levy MM, Miyasaki A, Langston D. Work of breathing as a weaning parameter in mechanically ventilated patients. *Chest* 1995;108(4):1018-1020.
 101. Kirton OC, DeHaven CB, Morgan JP, Windsor J, Civetta JM. Elevated imposed work of breathing masquerading as ventilator weaning intolerance. *Chest* 1995;108(4):1021-1025.
 102. Shikora SA, Bistrrian OR, Borlase BC, Blackburn GL, Stone MD, Benotti PN. Work of breathing: reliable predictor of weaning and extubation. *Crit Care Med* 1990;18(2):157-162.
 103. Harpin RP, Baker JP, Downer JP, Whitwell J, Gallacher WN. Correlation of the oxygen cost of breathing and length of weaning from mechanical ventilation. *Crit Care Med* 1987;15(9):807-812.
 104. Shikora SA, Benotti PN, Johannigman JA. The oxygen cost of breathing may predict weaning from mechanical ventilation better than the respiratory rate to tidal volume ratio. *Arch Surg* 1994;129(3):269-274.
 105. Tahvanainen J, Salmenpera M, Nikki P. Extubation criteria after weaning from intermittent mandatory ventilation and continuous positive airway pressure. *Crit Care Med* 1983;11(9):702-707.
 106. Hubble CL, Gentile MA, Tripp DS, Craig DM, Meliones JN, Cheifetz IM. Dead-space to tidal volume ratio predicts successful extubation in infants and children. *Crit Care Med* 2000;28(6):2034-2040.
 107. Capdevila XJ, Perrigault PF, Perey PJ, Roustan JPA, d'Athis F. Occlusion pressure and its ratio to maximum inspiratory pressure are useful predictors for successful extubation following T-piece weaning trial. *Chest* 1995;108(2):482-489.
 108. Montgomery AB, Holle RH, Neagley SR, Pierson DJ, Schoene RB. Prediction of successful ventilator weaning using airway occlusion pressure and hypercapnic challenge. *Chest* 1987;91(4):496-499.
 109. Gandia F, Blanco J. Evaluation of indexes predicting the outcome of ventilator weaning and value of adding supplemental inspiratory load. *Intensive Care Med* 1992;18(6):327-333.
 110. Gillespie LM, White SD, Sinha SK, Donn SM. Usefulness of the minute ventilation test in predicting successful extubation in newborn infants: a randomized controlled trial. *J Perinatol* 2003;23(3):205-207.
 111. Smina M, Salam A, Khamiees M, Gada P, Amoateng-Adjepong Y, Manthous CA. Cough peak flows and extubation outcomes. *Chest* 2003;124(1):262-268.

112. Bach JR, Saporito LR. Criteria for extubation and tracheostomy tube removal for patients with ventilatory failure: a different approach to weaning. *Chest* 1996;110(6):1566-1571.
113. Martinez A, Seymour C, Nam M. Minute ventilation recovery time: a predictor of extubation outcome. *Chest* 2003;123(4):1214-1221.
114. Bruton A. A pilot study to investigate any relationship between sustained maximal inspiratory pressure and extubation outcome. *Heart Lung* 2002;31(2):141-149.
115. Balsan MJ, Jones JG, Watchko JF, Guthrie RD. Measurements of pulmonary mechanics prior to the elective extubation of neonates. *Pediatr Pulmonol* 1990;9(4):238-243.
116. Davis PG, Henderson-Smart DJ. Extubation from low-rate intermittent positive airways pressure versus extubation after a trial of endotracheal continuous positive airways pressure in intubated preterm infants. *Cochrane Database Syst Rev* 2001;(4):CD001078.
117. Zeggwagh AA, Abouqal R, Madani N, Zekraoui A, Kerkeb O. Weaning from mechanical ventilation: a model for extubation. *Intensive Care Med* 1999;25(10):1077-1083.
118. Redmond JM, Greene PS, Goldsborough MA, Cameron DE, Stuart RS, Sussman MS, et al. Neurologic injury in cardiac surgical patients with a history of stroke. *Ann Thorac Surg* 1996;61(1):42-47.
119. Harel Y, Vardi A, Quigley R, Brink LW, Manning SC, Carmody TJ, Levin DL. Extubation failure due to post-extubation stridor is better correlated with neurologic impairment than with upper airway lesions in critically ill pediatric patients. *Int J Pediatr Otorhinolaryngol* 1997;39(2):147-158.
120. Coplin WM, Pierson DJ, Cooley KD, Newell DW, Rubenfeld GD. Implications of extubation delay in brain-injured patients meeting standard weaning criteria. *Am J Respir Crit Care Med* 2000;161(5):1530-1536.
121. Khamiees M, Raju P, DeGirolamo A, Amoateng-Adjepong Y, Manthous CA. Predictors of extubation outcome in patients who have successfully completed a spontaneous breathing trial. *Chest* 2001;120(4):1262-1270.
122. Deem S, Bishop MJ. Evaluation and management of the difficult airway. *Crit Care Clin* 1995;11(1):1-27.
123. American Society of Anesthesiologists. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology* 2003;98(5):1269-1277. *Erratum in Anesthesiology* 2004;101(2):565.
124. Cork R, Monk JE. Management of a suspected and unsuspected difficult laryngoscopy with the laryngeal mask airway. *J Clin Anesth* 1992;4(3):230-234.
125. Van Boven MJ, Lengele B, Fraselle B, Butera G, Veyckemans F. Unexpected difficult tracheal reintubation after thyroglossal duct surgery: functional imbalance aggravated by the presence of a hematoma. *Anesth Analg* 1996;82(2):423-425.
126. Mort TC. Extubating the difficult airway: determining the odds of failure: be alert to the signs that flag a management challenge. *J Crit Illn* 2003;18(4):177-184.
127. Loudermilk EP, Hartmannsgruber M, Stoltzfus DP, Langevin PB. A prospective study of the safety of tracheal extubation using a pediatric airway exchange catheter for patients with a known difficult airway. *Chest* 1997;111(6):1660-1665.
128. Hartmannsgruber MW, Loudermilk E, Stoltzfus D. Prolonged use of a cook airway exchange catheter obviated the need for postoperative tracheostomy in an adult patient. *J Clin Anesth* 1997;9(6):496-498.
129. Hartmannsgruber MW, Rosenbaum SH. Safer endotracheal tube exchange technique (letter). *Anesthesiology* 1998;88(6):1683.
130. Davis S, Worley S, Mee RB, Harrison AM. Factors associated with early extubation after cardiac surgery in young children. *Pediatr Crit Care Med* 2004;5(1):63-68.
131. Venkataraman ST, Khan N, Brown A. Validation of predictors of extubation success and failure in mechanically ventilated infants and children. *Crit Care Med* 2000;28(8):2991-2996.
132. Fisher MM, Raper RF. The 'cuff-leak' test for extubation. *Anaesthesia* 1992;47(1):10-12.
133. Potgieter PD, Hammond JM. "Cuff" test for safe extubation following laryngeal edema (letter). *Crit Care Med* 1988;16(8):818.
134. De Bast Y, De Backer D, Moraine JJ, Lemaire M, Vandenberght C, Vincent JL. The cuff leak test to predict failure of extubation for laryngeal edema. *Intensive Care Med* 2002;28(9):1267-1272.
135. Miller RL, Cole RP. Association between reduced cuff leak volume and postextubation stridor. *Chest* 1996;110(4):1035-1040.
136. Jaber S, Chanques G, Matecki S, Ramonatxo M, Vergne C, Souche B, et al. Post-extubation stridor in intensive care unit patients: risk factors evaluation and importance of the cuff-leak test. *Intensive Care Med* 2003;29(1):69-74.
137. Marik PE. The cuff-leak test as a predictor of postextubation stridor: a prospective study. *Respir Care* 1996;41(6):509-511.
138. Sandhu RS, Pasquale MD, Miller K, Wasser TE. Measurement of endotracheal tube cuff leak to predict postextubation stridor and need for reintubation. *J Am Coll Surg* 2000;190(6):682-687.
139. Engoren M. Evaluation of the cuff-leak test in a cardiac surgery population. *Chest* 1999;116(4):1029-1031.
140. Kemper KJ, Ritz RH, Benson MS, Bishop MS. Helium-oxygen mixture in the treatment of postextubation stridor in pediatric trauma patients. *Critical Care Med* 1991;19(3):356-359.
141. Adderley RJ, Mullins GC. When to extubate the croup patient: the "leak" test. *Can J Anaesth* 1987;34(3 Pt 1):304-306.
142. Seid AB, Godin MS, Pransky SM, Kearns DB, Peterson BM. The prognostic value of endotracheal tube-air leak following tracheal surgery in children. *Arch Oto-*

- laryngol Head Neck Surg 1991;117(8):880-882.
143. Morganroth ML, Morganroth JL, Nett LM, Petty TL. Criteria for weaning from prolonged mechanical ventilation. *Arch Intern Med* 1984;144(5):1012-1016.
 144. Clochesy JM, Daly BJ, Montenegro HD. Weaning chronically critically ill adults from mechanical ventilatory support: a descriptive study. *Am J Crit Care* 1995;4(2):93-99.
 145. Biery DR, Marks JD, Schapera A, Autry M, Schlobohm RM, Katz JA. Factors affecting perioperative pulmonary function in acute respiratory failure. *Chest* 1990;98(6):1455-1462.
 146. Hammond MD, Bauer KA, Sharp JT, Rocha RD. Respiratory muscle strength in congestive heart failure. *Chest* 1990;98(5):1091-1094.
 147. Rothaar RC, Epstein SK. Extubation failure: magnitude of the problem, impact on outcomes, and prevention. *Curr Opin Crit Care* 2003;9(1):59-66.
 148. Sapijaszko MJ, Brant R, Sandham D, Berthiaume Y. Nonrespiratory predictor of mechanical ventilation dependency in intensive care unit patients. *Crit Care Med* 1996;24(4):601-607.
 149. Smith IE, Shneerson JM. A progressive care programme for prolonged ventilatory failure: analysis of outcome. *Br J Anaesth* 1995;75(4):399-404.
 150. Scheinhorn DJ, Hassenpflug M, Artinian BM, LaBree L, Catlin JL. Predictors of weaning after 6 weeks of mechanical ventilation. *Chest* 1995;107(2):500-505.
 151. Cerra FB. Hypermetabolism, organ failure and metabolic support. *Surgery* 1987;101(1):1-14.
 152. Aubier M, Murciano D, Lecocguic Y, Viires N, Jacquens Y, Squara P, Pariente R. Effect of hypophosphatemia on diaphragmatic contractility in patients with acute respiratory failure. *N Engl J Med* 1985;313(7):420-424.
 153. Aubier M, Viires N, Piquet J, Murciano D, Blanchet F, Marty C, Gherardi R, Pariente R. Effects of hypocalcemia on diaphragmatic strength generation. *J Appl Physiol* 1985;58(7):2054-2061.
 154. Pingleton SK, Harmon GS. Nutritional management in acute respiratory failure. *JAMA* 1987;257(22):3094-3099.
 155. Lewis MI, Sieck GC, Fournier M, et al. The effect of nutritional deprivation of diaphragm contractility and muscle fiber size. *J Appl Physiol* 1986;60(2):596-603.
 156. Doekel RC Jr, Zwillich C, Scoggin CH, Kryger M, Weil JV. Clinical semi-starvation: depression of hypoxic ventilatory response. *N Engl J Med* 1976;295(7):358-361.
 157. Larca L, Greenbaum DM. Effectiveness of intensive nutritional regimes in patients who fail to wean from mechanical ventilation. *Crit Care Med* 1982;10(5):297-300.
 158. Bassili HR, Deitel M. Effect of nutritional support on weaning patients off mechanical ventilators. *J Parenter Enteral Nutr* 1981;5(2):161-163.
 159. Lyons KA, Brill RJ, Wieman RA, Jacobs BR. Continuation of transpyloric feeding during weaning of mechanical ventilation and tracheal extubation in children: a randomized controlled trial. *J Parenter Enteral Nutr* 2002;26(3):209-213.
 160. American Academy of Pediatrics Committee on Drugs: Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures. *Pediatrics* 1992;89(6 Pt 1):1110-1115.
 161. Gefke K, Andersen LW, Friesel E. Lidocaine given intravenously as a suppressant of cough and laryngospasm in connection with extubation after tonsillectomy. *Acta Anaesthesiol Scand* 1983;27(2):111-112.
 162. Staffler JG, Weissler MC, Tyler EP, Drake AF. The prevention of postoperative stridor and laryngospasm with topical lidocaine. *Arch Otolaryngol Head Neck Surg* 1991;117(10):1123-1128.
 163. Markovitz BP, Randolph AG. Corticosteroids for the prevention and treatment of postextubation stridor in neonates, children and adults. *Cochrane Database Syst Rev* 2000;(2):CD001000.
 164. Freezer N, Butt W, Phelan P. Steroids in croup: do they increase the incidence of successful extubation? *Anaesth Intensive Care* 1990;18(2):224-228.
 165. Tibballs J, Shann FA, Landau LI. Placebo-controlled trial of prednisolone in children intubated for croup. *Lancet* 1992;340(8822):745-748.
 166. Steer PA, Henderson-Smart DJ. Caffeine versus theophylline for apnea in preterm infants. *Cochrane Database Syst Rev* 2000;(2):CD000273.
 167. Henderson-Smart DJ, Davis PG. Prophylactic methylxanthines for extubation in preterm infants. *Cochrane Database Syst Rev* 2003;(1):CD000139.
 168. Listello D, Sessler CN. Unplanned extubation: clinical predictors for reintubation. *Chest* 1994;105(5):1496-1503.
 169. Epstein SK. Etiology of extubation failure and the predictive value of the rapid shallow breathing index. *Am J Respir Crit Care Med* 1995;152(2):545-549.
 170. Whelan J, Simpson SQ, Levy H. Unplanned extubation: predictors of successful termination of mechanical ventilatory support. *Chest* 1994;105(6):1808-1812.
 171. Tindol GA Jr, DiBenedetto RJ, Kosciuk L. Unplanned extubations. *Chest* 1994;105(6):1804-1807.
 172. Vassal T, Anh NG, Gabillet JM, Guidet B, Staikowsky F, Offenstadt G. Prospective evaluation of self-extubations in a medical intensive care unit. *Intensive Care Med* 1993;19(6):340-342.
 173. Franck LS, Vaughan B, Wallace J. Extubation and reintubation in the NICU: identifying opportunities to improve care. *Pediatr Nurs* 1992;18(3):267-270.
 174. De Paoli AG, Davis PG, Faber B, Morley CJ. Devices and pressure sources for administration of nasal continuous positive airway pressure (NCPAP) in preterm neonates. *Cochrane Database Syst Rev* 2002;(4):CD002977.
 175. Jiang JS, Kao SJ, Wang SN. Effect of early application of biphasic positive airway pressure on the outcome of extubation in ventilator weaning. *Respirology* 1999;4(2):161-165.
 176. Meyer TJ, Hill NS. Noninvasive positive pressure ven-

- tilation to treat respiratory failure. *Ann Intern Med* 1994;120(9):760-770.
177. Keenan SP, Powers C, McCormack DG, Block G. Noninvasive positive-pressure ventilation for postextubation respiratory distress: a randomized controlled trial. *JAMA* 2002;287(24):3238-3244.
 178. Kilger E, Briegel J, Haller M, Frey L, Schelling G, Stoll C, et al. Effects of noninvasive positive pressure ventilatory support in non-COPD patients with acute respiratory insufficiency after early extubation. *Intensive Care Med* 1999;25(12):1374-1380.
 179. Davies MW, Davis PG. Nebulized racemic epinephrine for extubation of newborn infants. *Cochrane Database Syst Rev* 2002;(1):CD000506.
 180. Liebler JM, Markin CJ. Fiberoptic bronchoscopy for diagnosis and treatment. *Crit Care Clin* 2000;16(1):83-100.
 181. Walker P, Forte V. Failed extubation in the neonatal intensive care unit. *Ann Otol Rhinol Laryngol* 1993;102(7):489-495.
 182. Plummer AL, Gracey DR. Consensus conference on artificial airways in patients receiving mechanical ventilation. *Chest* 1989;96(1):178-180.
 183. Maziak D, Meade M, Todd TR. The timing of tracheostomy: a systematic review. *Chest* 1998;114(2):605-609.
 184. Garner JS. Hospital Infection Control Practices Advisory Committee, Centers for Disease Control and Prevention. Guidelines for isolation precautions in hospitals. 1-01-1996. Available at www.cdc.gov.
 185. Centers for Disease Control & Prevention. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health-care facilities. *MMWR* 43(RR13);1-132. Publication date 10/28/1994. Available at www.cdc.gov.
 186. Lange JH. SARS respiratory protection (letter). *CMAJ* 2003;169(6):541-542.
 187. Ho PL, Tang XP, Seto WH. SARS: hospital infection control and admission strategies. *Respirology* 2003;8 Suppl:S41-S45.
 188. Tablan OC, Anderson LJ, Besser R, Bridges C, Hajjeh R. CDC. Healthcare Infection Control Practices Advisory Committee. Guidelines for preventing health-care-associated pneumonia, 2003: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. *MMWR Recommendations & Reports* 2004;53(RR-3):1-36.
 189. Bolyard EA, Tablan OC, Williams WW, Pearson ML, Shapiro CN, Deitchmann SD. Guideline for infection control in healthcare personnel, 1998. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1998;19(6):407-463. *Erratum in Infect Control Hosp Epidemiol* 1998;19(7):493.

Interested persons may photocopy these Guidelines for noncommercial purposes of scientific or educational advancement. Please credit AARC and RESPIRATORY CARE Journal.

All of the AARC CPGs may be downloaded at no charge from
<http://www.RCJournal.com/>