

# The Timed Inspiratory Effort Index as a Weaning Predictor: Analysis of Intra- and Interobserver Reproducibility

Arthur E da Silva Neto, Leonardo C de Souza, Helson LLS Costa, Bruno LS Guimarães, Leandro M de Azeredo, Marcos David P Godoy, and Jocemir R Lugon

**BACKGROUND:** Prolonged ventilatory weaning may expose patients to unnecessary discomfort, increase the risk of complications, and raise the costs of hospital treatment. In this scenario, indexes that reliably predict successful liberation can be helpful. **OBJECTIVE:** To evaluate the intra- and interobserver reproducibility of the timed inspiratory effort index as a weaning predictor. **METHODS:** This prospective observational study included subjects judged as able to start liberation from mechanical ventilation. For the intra-observer analysis, the same investigator performed 2 measurements in each selected patient with an interval of 30 min a rest. For interobserver analysis, 2 measurements were obtained in another sample of subjects, also with an interval of 30 min rest, but each of one performed by a different investigator. The Bland-Altman diagram, the coefficient concordance of kappa, and the Pearson correlation coefficient were used to compare the measurements. The performance of the timed inspiratory effort index was assessed by receiver operating characteristic curves. Values of  $P < .05$  were considered significant. **RESULTS:** We selected 113 subjects (43 males; mean  $\pm$  SD age,  $77 \pm 14$  y). Fifty-six (49.6%) achieved successful liberation, and 33 (29%) died in the ICU. The mean  $\pm$  SD duration of mechanical ventilation was  $14.4 \pm 6.7$  d. The Bland-Altman diagrams that addressed intra- and interobservers agreement showed low variability between measurements. Values of the concordance coefficients of kappa were 0.82 (0.68–0.95) and 0.80 (0.65–0.94), and of the linear correlation coefficients, 0.86 (0.77–0.91) and 0.89 (0.82–0.93) for the intra- and interobservers measurements, respectively. The mean  $\pm$  SD values for the area under the curve for each pair of the intra- and interobserver measurements were  $0.96 \pm 0.07$  versus  $0.94 \pm 0.07$  ( $P = .41$ ) and  $0.94 \pm 0.05$  versus  $0.90 \pm 0.07$  ( $P = .14$ ), respectively. **CONCLUSIONS:** The variability of the measurement of the timed inspiratory effort index by intra- and interobservers showed very high reproducibility, which reinforced the index as a sensible, accurate, and reliable outcome predictor of liberation from mechanical ventilation. *Key words:* ventilatory weaning; mechanical ventilation liberation; inspiratory muscles; variability; reproducibility; accuracy. [Respir Care 2020;65(5):636–642. © 2020 Daedalus Enterprises]

## Introduction

Mechanical ventilation is an essential therapy for patients with acute respiratory failure. Weaning is the process of transition from mechanical ventilation to spontaneous

breathing in patients who remained on mechanical ventilation for  $\geq 24$  h.<sup>1,2</sup> It should be emphasized that the withdrawal of the patient from mechanical ventilation may be more difficult than maintenance. Some researchers describe weaning as an obscure area of intensive care and

---

Drs da Silva Neto, Costa, Guimarães, de Azeredo, Godoy, and Lugon are affiliated with the Medical Science Post-graduation Program, Universidade Federal Fluminense, Niterói, Rio de Janeiro, Brazil. Dr de Souza is affiliated with the Physical Therapy School, Universidade Estácio de Sá, Niterói, Rio de Janeiro, Brazil. Drs da Silva Neto and de Souza are affiliated with the Intensive Care Division, Hospital Icaraí, Niterói, Rio de Janeiro, Brazil. Drs de Souza and Godoy are affiliated with the Intensive Care

---

Division, Hospital and Clínica São Gonçalo, São Gonçalo, Rio de Janeiro, Brazil. Dr Lugon is affiliated with the Medicine/Nephrology, Medical School of Universidade Federal Fluminense, Niterói, Rio de Janeiro, Brazil.

The authors have disclosed no conflicts of interest.

comment that, even with specialists, this stage of ventilatory care can be considered a mixture of art and science.<sup>2-4</sup>

The use of mechanical ventilation can cause ventilator-induced diaphragm dysfunction and generate changes in muscle structure and contractility, which results in reduced respiratory muscle strength, a process that can start as early as 18 h after mechanical ventilation.<sup>5-7</sup> Many researchers agree that the ventilator-induced diaphragm dysfunction can prolong weaning and is associated with an increased ICU stay and with hospital mortality.<sup>6,8-11</sup> In this regard, indexes that could evaluate for the presence of ventilator-induced diaphragm dysfunction and reliably predict a successful weaning would be helpful. The maximum inspiratory pressure ( $P_{I_{max}}$ ) was the first respiratory index described in the literature as a predictor of ventilatory liberation. Perhaps due to the lack of standardization in its measurement, the performance of  $P_{I_{max}}$  as a weaning predictor is poor in most studies.<sup>12-15</sup>

In the search for an index with better predictive power for mechanical ventilator liberation, we considered the combination of parameters (respiratory center stimulation and muscle response) that may be crucially impaired in patients on mechanical ventilation. The timed inspiratory effort index is a recently proposed weaning index that seems to be able to indicate readiness for mechanical ventilation liberation while providing a quick test of diaphragm strength, especially in patients on long-term mechanical ventilation associated with neurologic and neuromuscular diseases. Accordingly, the index showed better performance than those previously reported in a number of studies.<sup>16-18</sup> Thereby, the aim of this study was to evaluate the reliability of the timed inspiratory effort index as a weaning predictor by analyzing its intra- and interobserver variability and reproducibility.

## Methods

This was a prospective, observational study that involved subjects on mechanical ventilation admitted to the ICU of a general hospital from March 2017 to June 2018. The ethics committee of the Federal Fluminense University approved the study under the number CAAE 43049615.0000.5243. Informed consent was obtained from every patient, whenever possible, or from his or her next of kin. Enrollment

---

This study was partially supported by Universidade Estácio de Sá by using a research productivity program.

Correspondence: Leonardo Cordeiro de Souza PT PhD, Physiotherapy of Hospital Icaraí and Hospital and Clínica São Gonçalo, Rua Marquês de Paraná, 233 – 5o. andar. CEP 24030-215, Centro, Niterói, Rio de Janeiro, Brazil. E-mail: leonardo.cordeiro.souza@gmail.com.

DOI: 10.4187/respcare.07225

## QUICK LOOK

### Current knowledge

Premature discontinuation as well as unnecessary delay of liberation from mechanical ventilation have been associated with poor outcomes. Clinical judgment alone may not be enough for the guidance of the timing of withdrawal from mechanical ventilation. The timed inspiratory effort index combines the ability to generate an inspiratory pressure with the time needed to achieve the maximum inspiratory pressure.

### What this paper contributes to our knowledge

The high accuracy and reproducibility of the timed inspiratory effort index reinforce this weaning predictor as a reliable complementary tool for the clinician. This information may allow more specific programming of respiratory therapy for this population of critically ill patients.

criteria included patients >18 y of age, with an endotracheal or tracheostomy tube, who were judged as able to start the weaning process. They should also have resolution of the acute phase of the disease that led to mechanical ventilation and an adequate level of consciousness and cough reflex. Accordingly, subjects should have their infection under control, stable cardiovascular status (heart rate  $\leq 120$  beats/min and systolic blood pressure between 90 and 160 mm Hg, no or minimal use of vasopressors), stable metabolic status, hemoglobin  $> 10$  g/dL, adequate oxygenation (arterial oxygen saturation  $> 90\%$  with  $F_{IO_2} \leq 0.4$  or  $P_{aO_2}/F_{IO_2} \geq 150$ ) with final positive expiratory pressure  $\leq 8$  cm  $H_2O$ ), adequate frequency ( $\leq 35$  breaths/min), pressure support  $\leq 20$  cm  $H_2O$ , and no significant respiratory acidosis (pH  $> 7.30$ ). We excluded patients with tracheal stenosis, intracranial pressure  $> 20$  mm Hg, on sedation, with severe cardiac insufficiency or hemodynamic instability, signs of systemic infection or reinfection during the liberation process, or discontinuation due to other complications.

## Procedures and Definitions

All the subjects in the study were on mechanical ventilation with Extend XT (TAEMA, Air Liquide, Paris, France) and were continuously monitored by using the DX 2010 multi-parameter monitor (Dixtal, São Paulo, São Paulo, Brazil), which records the electrocardiogram, heart rate,  $S_{pO_2}$ , and systemic arterial pressure. At any sign of instability, the test was interrupted and the subject was returned to mechanical ventilation. For measurements of  $P_{I_{max}}$  and the




Fig. 1. Typical record, which demonstrates that the maximum inspiratory peak pressure (61 cm H<sub>2</sub>O, vertical dashed line) over 60 s of observation was achieved at 58.2 s. In this test, the timed inspiratory effort index was 1.04 cm H<sub>2</sub>O/s.

timed inspiratory effort index, the digital vacuumeter (Magnamed, São Paulo, São Paulo, Brazil), with a scale of 300 cm H<sub>2</sub>O and an increment of 0.1 cm H<sub>2</sub>O and a time interval of 100 ms, was used. The method used was the occlusion of the airway during inspiration with a unidirectional valve,<sup>19,20</sup> with the subjects positioned in the supine position, and the head elevated at 45°. The cuff was hyperinflated to prevent air leak during measurement. After tracheal aspiration, the subjects remained connected to the mechanical ventilator to rest for 2 min on 100% oxygen.<sup>13,14,16-18</sup> After hyperoxygenation, the ventilator was disconnected, and the subject breathed spontaneously. After 10 s of spontaneous breathing, the digital manometer was connected to the artificial airway at the end of a normal expiration (at FRC). The airway was occluded for 60 s while the values were recorded, which corresponded to each inspiratory effort.<sup>13,14,16-18</sup>

For the intra-observer analysis, the same respiratory physiotherapist performed 2 measurements in each selected subject, with a 30-min rest period on pressure-support ventilation. For interobserver analysis, 2 measurements were obtained by different respiratory physiotherapists on another group of subjects, which also provided a 30-min rest period between tests. The timed inspiratory effort index was calculated as the ratio of the  $P_{I_{max}}$  registered after the first 30 s of observation to the corresponding time to reach this value of  $P_{I_{max}}$ , while keeping the airway occlusion with a unidirectional valve for up to 60 s. The timed inspiratory effort index =  $P_{I_{max}}$ (after 30 s of start, cm H<sub>2</sub>O)/time (in s) to achieve this value of  $P_{I_{max}}$ . A typical record of inspiratory pressures by time is shown in Figure 1.

Liberation was considered successful if spontaneous breathing was maintained for >48 h after mechanical

ventilation withdrawal. The decision to return to mechanical ventilation was taken by a respiratory physiotherapist and the attending physician (both did not have access to the results of the timed inspiratory effort index measurement). Criteria to interrupt periods of spontaneous breathing during the T-piece trial were as follows: agitation and/or anxiety, reduced level of consciousness, diaphoresis, dyspnea, and/or cyanosis, oxygen saturation measured from an arterial blood sample < 90% or  $P_{aO_2}$  < 60 mm Hg, with  $F_{IO_2}$  > 0.4,  $P_{aCO_2}$  > 50 mm Hg or increased in >8 mm Hg, arterial pH < 7.33 or decreased by  $\geq 0.07$ , frequency > 35 breaths/min or increased by 50% for  $\geq 5$  min, heart rate > 140 beats/min or a sustained increase or decrease > 20%, mean blood pressure > 130 mm Hg or < 70 mm Hg, and tachyarrhythmias.<sup>2,12-14,16,17,18,21</sup>

### Statistical Analysis

The sample size was calculated to be able to evaluate the differences in the order of 20% between groups with a power of 80% and an alpha error of 5%, in which a minimum number of 35 individuals was found for each studied group. The results were expressed as mean  $\pm$  SD for continuous variables with normal distribution and, alternatively, as median and 95% CI. Category variables were expressed as frequencies. The differences between the continuous variables and the frequencies were evaluated by using the Wilcoxon test and the chi-square test, respectively.

The obtained values of the timed inspiratory effort index were evaluated as to their ability to predict weaning by calculating the area under the receiver operating characteristic curve. The areas under the receiver operating characteristic curve were compared by using the method of Hanley and

# TIMED INSPIRATORY EFFORT INDEX AS A WEANING PREDICTOR

Table 1. Subject Characteristics based on Weaning Outcome

Variable	Total	Weaning		P (Success vs Failure)
		Success	Failure	
Sample, <i>n</i> (%)	113	56 (49.6)	57 (50.4)	NA
Males/females, <i>n</i>	43/70	17/39	26/31	.14
Race (White/Black), <i>n</i>	67/46	34/18	33/28	.87
Age, mean ± SD y	77 ± 14	75.2 ± 15.3	78.6 ± 13.6	.21
APACHE II score, mean ± SD	18.9 ± 5.5	17.7 ± 5.0	20 ± 5.9	.03
Duration of MV, mean ± SD d	14.4 ± 6.7	12.8 ± 6.4	16.1 ± 6.6	<.01
P <sub>aCO2</sub> before test, mean ± SD mm Hg	40.9 ± 4.3	40.2 ± 3.9	41.7 ± 4.7	.07
Deaths in the ICU, <i>n</i> (%)	33 (29.2)	12 (10.6)	21 (18.6)	.07
Tracheostomy, <i>n</i> (%)	63 (55.8)	24 (21.3)	39 (34.5)	<.01

NA = not applicable

APACHE = Acute Physiology and Chronic Health Evaluation

MV = mechanical ventilation

Table 2. Values Obtained with the Same Observer (Group Intra-observation, *n* = 60)

Variable	Examination, Median (95% CI)		P
	First	Second	
P <sub>I<sub>max</sub></sub> , cm H <sub>2</sub> O	44.5 (36.8–49)	44 (38.8–48)	.13
Time to achieve P <sub>I<sub>max</sub></sub> , s	49.9 (47.5–53.6)	48.5 (43.1–53)	.30
Timed inspiratory effort index	0.87 (0.73–1.01)	0.85 (0.79–1.01)	.16

P<sub>I<sub>max</sub></sub> = maximum inspiratory pressure

McNeil.<sup>22</sup> Specifically for the analysis of intra- and interobserver reproducibility, the Bland-Altman diagram and the inter-rater agreement statistic (kappa) were used. Values of *P* < .05 were considered significant. Statistical analysis was performed by using the statistical program SPSS, version 18.0 (Chicago, Illinois) and the MedCalc statistical program, version 11.4.2.0. (Mariakerke, Oost-Vlaanderen, Belgium).

## Results

One hundred and thirteen subjects were selected (43 males, mean ± SD age, 77 ± 14 y). Fifty-six (49.6%) underwent successful liberation, and 33 (29%) died in the ICU. Tracheotomy was necessary in 63 subjects (55.8%), the mean ± SD duration of mechanical ventilation was 14.4 ± 6.7 d, and the mean ± SD APACHE (Acute Physiology and Chronic Health Evaluation) II score was 18.9 ± 5.5 (Table 1). The median and 95% CI of the intra-observer examinations with regard to P<sub>I<sub>max</sub></sub>, time to achieve P<sub>I<sub>max</sub></sub>, and timed inspiratory effort index are depicted in Table 2. In this setting, no statistically significant difference was found between measurements: *P* = .13 for P<sub>I<sub>max</sub></sub>, *P* = .30 for the time to reach P<sub>I<sub>max</sub></sub>, and *P* = .16 for the timed

Table 3. Values Obtained with Different Observers (Group Interobservation, *n* = 53)

Variable	Second Observer	Third Observer	P
P <sub>I<sub>max</sub></sub> , cm H <sub>2</sub> O	49 (44–55.1)	51 (43.8–55.1)	.30
Time to achieve P <sub>I<sub>max</sub></sub> , s	48.4 (42.3–53.6)	49 (45.8–54.3)	.99
Timed inspiratory effort index	1.02 (0.88–1.16)	1.02 (0.88–1.13)	.33

Data are median (95% CI).

P<sub>I<sub>max</sub></sub> = maximum inspiratory pressure

inspiratory effort index. Correspondent results for the interobserver examinations are in Table 3. Again, statistically significant differences were not found, with *P* = .30 for P<sub>I<sub>max</sub></sub>, *P* = .99 for the time to reach P<sub>I<sub>max</sub></sub>, and *P* = .33 for the timed inspiratory effort index.

The Bland-Altman diagrams that addressed intra- and interobserver agreement for these variables are in Figure 2. Values for the concordance coefficients of kappa of the intra- and interobservers were median weighted Kappa (95% CI) 0.82 (0.68–0.95) and 0.80 (0.65–0.94), respectively. Similar findings were found for linear correlation coefficients of the intra- and interobservers r<sup>2</sup> (95% CI) linear regression of Pearson 0.86 (0.77–0.91) and 0.89 (0.82–0.93), respectively (Fig. 3). When evaluating the weaning predictive power of the measurements of the intra-observer data by receiver operating characteristic curves, no statistically significant difference was found between the curves (0.96 ± 0.07 vs 0.94 ± 0.07; *P* = .41); the analysis of the interobserver data showed comparable results (0.94 ± 0.05 vs 0.90 ± 0.07; *P* = .14) (Fig. 4).

## Discussion

P<sub>I<sub>max</sub></sub> was recommended as a weaning predictor by an International Consensus Conference<sup>2</sup>; however, even

## TIMED INSPIRATORY EFFORT INDEX AS A WEANING PREDICTOR




Fig. 2. A Bland-Altman diagram of the intra- (A) and interobservers (B). TIE = timed inspiratory effort. Center lines show means, and dotted lines denote  $\pm 1.96$  SD.




Fig. 3. Linear Pearson correlation of the intra- (A) and interobservers (B). TIE = timed inspiratory effort.




Fig. 4. The area under the receiver operating characteristic curve, evaluating the TIE index as a predictor of weaning success of the intra- (A) and interobservers (B). TIE = timed inspiratory effort.

when measured by adopting the unidirectional inspiratory valve, the index performs below expectation in this regard.<sup>12-15,17,19,20</sup> The timed inspiratory effort index is a new weaning index, easy to perform at bedside and that has been reported to be more accurate.<sup>14,17,18</sup> The aim of the present study was to assess the reliability and reproducibility of the timed inspiratory effort index to better validate the measure as a useful weaning predictor. For

this purpose, we resorted to a traditional analysis of intra- and interobserver measurements. The studied sample consisted of older subjects, with a predominance of females. They had a high APACHE II score, prolonged duration of mechanical ventilation, and a high rate of tracheostomy. In this scenario, the mortality rate (29.2%) cannot be seen as a surprise. The features of our subjects were common to the ones found in most of the ICUs worldwide. Half of

the subjects underwent successful liberation, a number also consistent with previous studies.<sup>10,14,18,23</sup>

As an initial approach to accomplish the study purpose, the median of the 2 intra-observer measurements were compared, and no statistical significant difference was found. The same occurred with the 2 measurements of the interobserver were compared. These findings demonstrated that the timed inspiratory effort values may be easily and reproducibly obtained. The graphic analysis of the Bland-Altman diagrams showed that the mean line of either intra- or the interobserver data crossed the y axis close to the zero value. In addition, most of the values in both graphics were within the limits of  $\pm 1.96$  SD. The kappa concordance coefficients were 0.82 and 0.80 for the intra- and interobservers, respectively, which denoted very good agreement between measurements. Accordingly, the linear correlation coefficients were 0.86 and 0.89 for the intra- and interobservers, which indicated a high correlation between measurements.

Historically, few weaning predictors were recommended for clinical practice, mainly due to their low accuracy and reproducibility. In previous studies, the values for the accuracy of the timed inspiratory effort index (referred to as the area under the receiver operating characteristic curve) ranged from 0.90 to 0.96, which expressed a better performance than other predictive indexes described in the literature.<sup>14,17,18</sup> Interestingly, the values for the areas under the curve in the present study also ranged from 0.90 to 0.96. More importantly, no statistical difference was observed between the curves of each of the 2 sets of measurements for the intra- or interobserver data. Given the nature of the study, design limitations were restricted to those that pertained to the pre-established methods used to analyze intra- and interobserver reproducibility. However, it should be commented that it seemed prudent to preclude measurements of the timed inspiratory effort index in circumstances such as elevated intracranial pressure, risk of abdominal content evisceration, acute coronary artery disease, flail chest, and unstable heart failure.

### Conclusions

The results of the present study, which addressed the variability of the measurement of the timed inspiratory effort index by intra- and interobservers, showed a very high reproducibility, which reinforced the index as a sensible, accurate, and reliable outcome predictor with regard to liberation from mechanical ventilation. Accordingly, analysis of our data suggested that a measure might be sufficient to make a decision in clinical settings.

### ACKNOWLEDGMENTS

The authors thank the respiratory physiotherapists and physicians of the ICU of the Hospital Icaraf, especially André Luiz da Cunha Serejo, Vitor

Ricardo Stanizio Daher, Leandro Possidente Daher, Raphaela Cristine Carvalho Cordeiro, and Wanderlei Augusto Junior.

### REFERENCES

- Béduneau G, Pham T, Schortgen F, Piquilloud L, Zogheib E, Jonas M, et al; WIND (Weaning according to a New Definition) Study Group and the REVA (Réseau Européen de Recherche en Ventilation Artificielle) Network. Epidemiology of weaning outcome according to a new definition: the WIND Study. *Am J Respir Crit Care Med* 2017;195(6):772-783.
- Boles JM, Bion J, Connors A, Herridge M, Marsh B, Melot C, et al. Weaning from mechanical ventilation. *Eur Respir J* 2007;29(5):1033-1056.
- Milic-Emili J. Is weaning an art or a science? *Am Rev Respir Dis* 1986;134(6):1107-1108.
- Barbas CS, Ísola AM, Farias AM, Cavalcanti AB, Gama AM, Duarte AC, et al. Brazilian recommendations of mechanical ventilation 2013. Part 1. *Rev Bras Ter Intensiva* 2014;26(3):215-239.
- Levine S, Nguyen T, Taylor N, Friscia ME, Budak MT, Rothenberg P, et al. Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. *N Engl J Med* 2008;358(13):1327-1335.
- Jaber S, Petrof BJ, Jung B, Chanques G, Berthet JP, Rabuel C, et al. Rapidly progressive diaphragmatic weakness and injury during mechanical ventilation in humans. *Am J Respir Crit Care Med* 2011;183(3):364-371.
- Powers SK, Kavazis AN, Levine S. Prolonged mechanical ventilation alters diaphragmatic structure and function. *Crit Care Med* 2009;37(10 Suppl):S347-S353.
- Demoule A, Jung B, Prodanovic H, Molinari N, Chanques G, Coirault C, et al. Diaphragm dysfunction on admission to the intensive care unit. Prevalence, risk factors, and prognostic impact—a prospective study. *Am J Respir Crit Care Med* 2013;188(2):213-219.
- Supinski GS, Callahan LA. Diaphragm weakness in mechanically ventilated critically ill subjects. *Crit Care* 2013;17(3):R120.
- Medrinal C, Prieur G, Frenoy É, Robledo Quesada AR, Poncet A, Bonnevie T, et al. Respiratory weakness after mechanical ventilation is associated with one-year mortality a prospective study. *Crit Care* 2016;20(1):231.
- De Jonghe B, Bastuji-Garin S, Durand MC, Malissin I, Rodrigues P, Cerf C, et al; Groupe de Réflexion et d'Etude des Neuromyopathies en Réanimation. Respiratory weakness is associated with limb weakness and delayed weaning in critical illness. *Crit Care Med* 2007;35(9):2007-2015.
- Nemer SN, Barbas CS, Caldeira JB, Cárias TC, Santos RG, Almeida LC, et al. A new integrative weaning index of discontinuation from mechanical ventilation. *Crit Care* 2009;13(5):R152.
- de Souza LC, Silva CTJr, Almeida JR, Lugon JR. Comparison of maximal inspiratory pressure, tracheal airway occlusion pressure, and its ratio in the prediction of weaning outcome: impact of the use of a digital vacuumeter and the unidirectional valve. *Respir Care* 2012;57(8):1285-1290.
- de Souza LC, Guimarães FS, Lugon JR. Evaluation of a new index of mechanical ventilation weaning: the timed inspiratory effort. *J Intensive Care Med* 2015;30(1):37-43.
- Sassoon CS, Mahutte CK. Airway occlusion pressure and breathing pattern as predictors of weaning outcome. *Am Rev Respir Dis* 1993;148(4 Pt 1):860-866.
- de Souza LC, da Silva CTJr, Lugon JR. Evaluation of the inspiratory pressure using a digital vacuumeter in mechanically ventilated subjects: analysis of the time to achieve the inspiratory peak. *Respir Care* 2012;57(2):257-262.
- de Souza LC, Guimarães FS, Lugon JR. The timed inspiratory effort: a promising index of mechanical ventilation weaning for subjects with

- neurologic or neuromuscular diseases. *Respir Care* 2015;60(2):231-238.
18. da Silva Guimarães BL, de Souza LC, Guimarães FS, Lugon JR. Serial weekly measurements of the timed inspiratory effort index can predict successful prolonged weaning. *Respir Care* 2019;64(10):1286-1292.
  19. Marini JJ, Smith TC, Lamb V. Estimation of inspiratory muscle strength in mechanically ventilated subjects: the measurement of maximal inspiratory pressure. *J Crit Care* 1986;1(1):32-38.
  20. Caruso P, Friedrich C, Denari SD, Ruiz SA, Deheinzeln D. The unidirectional valve is the best method to determine maximal inspiratory pressure during weaning. *Chest* 1999;115(4):1096-1101.
  21. Yang KL, Tobin MJ. A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991;324(21):1445-1450.
  22. Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology* 1983;148(3):839-843.
  23. Jubran A, Grant BJ, Duffner LA, Collins EG, Lanuza DM, Hoffman LA, Tobin MJ. Effect of pressure support vs unassisted breathing through a tracheostomy collar on weaning duration in subjects requiring prolonged mechanical ventilation: a randomized trial. *JAMA* 2013;309(7):671-677.